

MAXIMUM ENTROPY DISTRIBUTION OF ORDER STATISTICS WITH GIVEN MARGINALS

CRISTINA BUTUCEA, JEAN-FRANÇOIS DELMAS, ANNE DUTFOY, AND RICHARD FISCHER

ABSTRACT. We consider distributions of ordered random vectors with given one-dimensional marginal distributions. We give an elementary necessary and sufficient condition for the existence of such a distribution with finite entropy. In this case, we give explicitly the density of the unique distribution which achieves the maximal entropy and compute the value of its entropy. This density is the unique one which has a product form on its support and the given one-dimensional marginals. The proof relies on the study of copulas with given one-dimensional marginal distributions for its order statistics.

1. INTRODUCTION

Order statistics, an almost surely non-decreasing sequence of random variables, have received a lot of attention due to the diversity of possible applications. If $X = (X_1, \dots, X_d)$ is a d -dimensional random vector, then its order statistics $X^{OS} = (X_{(1)}, \dots, X_{(d)})$ corresponds to the permutation of the components of X in the non-decreasing order, so that $X_{(1)} \leq X_{(2)} \leq \dots \leq X_{(d)}$. The components of the underlying random vector X are usually, but not necessarily, independent and identically distributed (i.i.d.). Special attention has been given to extreme values $X_{(1)}$ and $X_{(d)}$, the range $X_{(d)} - X_{(1)}$, or the median value. Direct application of the distribution of the k -th largest order statistic occurs in various fields, such as climatology, extreme events, reliability, insurance, financial mathematics. We refer to the monographs of David and Nagaraja [7] and Arnold, Balakrishnan, and Nagaraja [1] for a general overview on the subject of order statistics. We are interested in the dependence structure of order statistics, which has received great attention when the underlying random vector is i.i.d. and for the non i.i.d. case as well. In the i.i.d. case, Bickel [3] showed that any two order statistics are positively correlated. The copula of the joint distribution of $X_{(1)}$ and $X_{(d)}$ is derived in Schmitz [18] with exact formulas for Kendall's τ and Spearman's ρ . In Avérous, Genest, and Kocher [2], it is shown that the dependence of the j -th order statistic on the i -th order statistic decreases as the distance between i and j increases according to the bivariate monotone regression dependence ordering. The copula connecting the limit distribution of the two largest order statistics, called bi-extremal copula, is given by de Melo Mendes and Sanfins [8] with some additional properties. Exact expressions for Pearson's correlation coefficient, Kendall's τ and Spearman's ρ for any two order statistics are obtained in Navarro and Balakrishnan [15]. For the non i.i.d. case, Kim and David [13] shows that some pairs of order statistics can be negatively correlated, if the underlying random vector is sufficiently negatively dependent. Positive dependence measures for two order statistics are considered in Boland, Hollander, Joag-Dev, and Kocher [4] when the underlying random variables are

Date: September 8, 2015.

2010 Mathematics Subject Classification. 62H05, 60E15, 62G30, 94A17.

Key words and phrases. copula, entropy, maximum entropy, order statistics.

This work is partially supported by the French "Agence Nationale de la Recherche", CIFRE n° 1531/2012, and by EDF Research & Development, Industrial Risk Management Department.

independent but arbitrarily distributed or when they are identically distributed but not independent. A generalization of these results for multivariate dependence properties is given by Hu and Chen [10]. See also Dubhashi and Häggström [9] for conditional distribution of order statistics.

Here, we focus on the cumulative distribution function (cdf) of order statistics without referring to an underlying distribution. That is, we consider random vectors $X = (X_1, \dots, X_d) \in \mathbb{R}^d$ such that a.s. $X_1 \leq \dots \leq X_d$ and we suppose that the one-dimensional marginal distributions $\mathbf{F} = (\mathbf{F}_i, 1 \leq i \leq d)$ are given, where $\mathbf{F}_i$ is the cdf of X_i . A necessary and sufficient condition for the existence of a joint distribution of order statistics with one-dimensional marginals $\mathbf{F}$ is that they are stochastically ordered, that is:

$$(1) \quad \mathbf{F}_{i-1}(x) \geq \mathbf{F}_i(x) \quad \text{for all } 2 \leq i \leq d, x \in \mathbb{R}.$$

With the marginals fixed, the joint distribution of the order statistics can be characterized by the connecting copula of the random vector, which contains all information on the dependence structure of the order statistics. Copulas of order statistics derived from an underlying i.i.d. sample were considered in [2] in order to calculate measures of concordance between any two pairs of order statistics. For order statistics derived from a general parent distribution, Navarro and Spizzichino [16] shows that the copula of the order statistics depends on the marginals and the copula of the parent distribution through an exchangeable copula and the average of the marginals. Construction of some copula of order statistics with given marginals were given in Lebrun and Dutfoy [14].

Our aim is to find the cdf of order statistics of dimension d with fixed marginals which maximizes the differential entropy H defined as, for a cdf F with density f :

$$H(F) = - \int f \log f,$$

and $H(F) = -\infty$ if F does not have a density. If Z is a random variable with cdf F , we shall use the convention $H(Z) = H(F)$. In an information-theoretic interpretation, the maximum entropy distribution is the least informative among order statistics with given marginals. This problem appears in models where the one-dimensional marginals are well known (either from different experimentation or from physical models) but the dependence structure is unknown. In Butucea, Delmas, Dutfoy, and Fischer [6], we gave, when it exists, the maximum entropy distribution of $(X_1, \dots, X_d)$ such that X_i is uniformly distributed on $[0, 1]$ for $1 \leq i \leq d$ and the distribution of $X_{(d)} = \max_{1 \leq i \leq d} X_i$ is given.

For a d -dimensional random variable $X = (X_1, \dots, X_d)$ with cdf F and copula C_F , the entropy of F can be decomposed into the sum of the entropy of its one-dimensional marginals plus the entropy of C_F (see Zhao and Lin [19]):

$$H(F) = \sum_{i=1}^d H(\mathbf{F}_i) + H(C_F),$$

where $\mathbf{F}_i$ is the cdf of X_i . In our case, since the marginals $\mathbf{F} = (\mathbf{F}_i, 1 \leq i \leq d)$ are fixed, maximizing the entropy of the joint distribution F of an order statistics is equivalent to maximizing the entropy of its copula C_F . Therefore we shall find the maximum entropy copula for order statistics with fixed marginal distributions. To solve this question, we introduce the

functional of $\mathbf{F}$ satisfying (1):

$$\mathbb{J}(\mathbf{F}) = \sum_{i=2}^d \int_{\mathbb{R}} \mathbf{F}_i(dt) |\log(\mathbf{F}_{i-1}(t) - \mathbf{F}_i(t))|.$$

The main result of this paper is given by Theorem 5.4 which we reproduce here.

Theorem. *Let $\mathbf{F} = (\mathbf{F}_i, 1 \leq i \leq d)$ be a d -dimensional vector of cdf's on $\mathbb{R}$ satisfying (1).*

- *If there exists $1 \leq i \leq d$ such that $H(\mathbf{F}_i) = -\infty$, or if $\mathbb{J}(\mathbf{F}) = +\infty$, then for all cdf F of order statistic with one-dimensional marginals $\mathbf{F}$, we have $H(F) = -\infty$.*
- *If $H(\mathbf{F}_i) > -\infty$ for all $1 \leq i \leq d$, and $\mathbb{J}(\mathbf{F}) < +\infty$, then there exists a unique cdf $F_{\mathbf{F}}$ of an order statistic with one-dimensional marginals $\mathbf{F}$ such that $H(F_{\mathbf{F}}) > -\infty$ and $H(F_{\mathbf{F}}) \geq H(F)$ for any cdf F of order statistic with one-dimensional marginals $\mathbf{F}$. Furthermore we have:*

$$H(F_{\mathbf{F}}) = d - 1 + \sum_{i=1}^d H(\mathbf{F}_i) - \mathbb{J}(\mathbf{F}),$$

and $F_{\mathbf{F}}$ has density $f_{\mathbf{F}}$ given for $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ by:

$$f_{\mathbf{F}}(x) = \mathbf{f}_1(x_1) \prod_{i=2}^d \frac{\mathbf{f}_i(x_i)}{\mathbf{F}_{i-1}(x_i) - \mathbf{F}_i(x_i)} \exp\left(-\int_{x_{i-1}}^{x_i} \frac{\mathbf{f}_i(s)}{\mathbf{F}_{i-1}(s) - \mathbf{F}_i(s)} ds\right) \mathbf{1}_{L^{\mathbf{F}}}(x),$$

where $\mathbf{f}_i$ is the density function of $\mathbf{F}_i$ and $L^{\mathbf{F}} \subset \mathbb{R}^d$ is the set of ordered vectors $(x_1, \dots, x_d)$, that is $x_1 \leq \dots \leq x_d$, such that $\mathbf{F}_{i-1}(t) > \mathbf{F}_i(t)$ for all $t \in (x_{i-1}, x_i)$ and $2 \leq i \leq d$. (Notice that this last condition is automatically fulfilled if $\mathbf{F}_{i-1} > \mathbf{F}_i$.)

The function $f_{\mathbf{F}}$ may be well defined even if $\mathbb{J}(\mathbf{F}) = +\infty$ and it might even be a density, see Remark 5.5. However in this case the entropy of the corresponding cdf is infinite. The density $f_{\mathbf{F}}$ has a product form on the domain $L^{\mathbf{F}}$: $f_{\mathbf{F}}(x) = \prod_{i=1}^d p_i(x_i) \mathbf{1}_{L^{\mathbf{F}}}(x)$ for $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ and some measurable functions p_i . Corollary 5.7 asserts that $f_{\mathbf{F}}$ is the only density which has a product form on the domain $L^{\mathbf{F}}$ and whose one-dimensional marginals are given by $\mathbf{F}$. This characterization will be used in a forthcoming paper on nonparametric estimation of $f_{\mathbf{F}}$.

Example. We consider the following example. Let $+\infty > \lambda_1 > \dots > \lambda_d > 0$ and for $1 \leq i \leq d$ let $\mathbf{F}_i$ be the cdf of the exponential distribution with mean $1/\lambda_i$ and density $\mathbf{f}_i(t) = \lambda_i e^{-\lambda_i t} \mathbf{1}_{\{t>0\}}$. Notice that $\mathbf{F}_{i-1} > \mathbf{F}_i$ on $(0, +\infty)$, so that $L^{\mathbf{F}} = \{(x_1, \dots, x_d) \in \mathbb{R}^d; 0 \leq x_1 \leq \dots \leq x_d\}$. It is easy to check that $\mathbb{J}(\mathbf{F}) < +\infty$ with $\mathbf{F} = (\mathbf{F}_i, 1 \leq i \leq d)$. Elementary computations yield that the maximum entropy density of the order statistic $(X_1, \dots, X_d)$, where X_i has distribution $\mathbf{F}_i$, is given by:

$$f_{\mathbf{F}}(x_1, \dots, x_d) = \mathbf{1}_{L^{\mathbf{F}}}(x) \lambda_1 e^{-\Delta_2 x_1} (1 - e^{-\Delta_2 x_1})^{\lambda_2/\Delta_2} \prod_{i=2}^d \lambda_i e^{-\Delta_{i+1} x_i} \frac{(1 - e^{-\Delta_{i+1} x_i})^{\lambda_{i+1}/\Delta_{i+1}}}{(1 - e^{-\Delta_i x_i})^{\lambda_{i-1}/\Delta_i}},$$

where $\Delta_i = \lambda_{i-1} - \lambda_i$ for $1 \leq i \leq d+1$ and $\lambda_{d+1} = 0$.

In the particular case $\lambda_i = (d-i+1)\lambda$ for some $\lambda > 0$, we get:

$$f_{\mathbf{F}}(x_1, \dots, x_d) = \mathbf{1}_{L^{\mathbf{F}}}(x) d! \lambda^d e^{-\lambda x_1} (1 - e^{-\lambda x_1})^{d-1} \prod_{i=2}^d \frac{e^{-\lambda x_i}}{(1 - e^{-\lambda x_i})^2}.$$

By considering the change of variable $u_i = 1 - e^{-\lambda x_i}$, we get the following result. For $1 \leq i \leq d$ let $\mathbf{F}_i$ be the cdf of the $\beta(1, d-i+1)$ distribution with density $\mathbf{f}_i(t) = (d-i+1)$

$1)(1-t)^{d-i}\mathbf{1}_{(0,1)}(t)$. Notice that $\mathbf{F}_{i-1} > \mathbf{F}_i$ on $(0,1)$. The maximum entropy density of the order statistic $(U_1, \dots, U_d)$, where U_i has distribution $\mathbf{F}_i$, is given by:

$$f_{\mathbf{F}}(u_1, \dots, u_d) = \mathbf{1}_{\{0 < u_1 < \dots < u_d < 1\}} d! u_1^{d-1} \prod_{i=2}^d \frac{1}{u_i^2}.$$

Elementary computations give $H(F_{\mathbf{F}}) = -\log(d!) + 2d - (d+1) \sum_{i=1}^d (1/i)$.

In order to prove Theorem 5.4, we first introduce the multidagonal of a copula. For a copula C on $\mathbb{R}^d$ and a random vector $U = (U_1, \dots, U_d)$ with cdf C , we consider its order statistics $U^{OS} = (U_{(1)}, \dots, U_{(d)})$ and define $\delta_{(i)}$ the one-dimensional cdf of $U_{(i)}$ for $1 \leq i \leq d$. Then the multidagonal of C is defined as $\delta_C = (\delta_{(i)}, 1 \leq i \leq d)$. This provides a generalization of the diagonal section of the copula C which corresponds to the cdf $\delta_{(d)}$ of $U_{(d)} = \max_{1 \leq i \leq d} U_i$. The maximum entropy copula with fixed diagonal section is given in [6]. The necessary and sufficient condition for a d -tuple δ to be the multidagonal of a (absolutely continuous) copula is provided by Jaworski and Rychlik [12]. In order to prove Theorem 5.4, we first establish a one-to-one map between the set of copulas of order statistics with given marginals $\mathbf{F}$ and symmetric copulas with a fixed multidagonal $\delta^{\mathbf{F}}$, which only depends on $\mathbf{F}$, see Lemma 3.4 and Proposition 3.13. Furthermore, this map preserves the absolute continuity of the copula, as well as the entropy up to an additive constant, which depends only on the fixed one-dimensional marginals $\mathbf{F}$, see Proposition 3.21. Then, under a necessary and sufficient condition on the multidagonal δ , we find the maximum entropy copula with a given general multidagonal δ and provide an explicit formula for its density, see Theorem 4.7. (Notice Theorem 4.7, which is first established to prove Theorem 5.4, can in fact be seen as a special case of Theorem 5.4.) The proof relies on the theory of optimization under infinite dimensional constraints developed by Borwein, Lewis, and Nussbaum [5].

The rest of the paper is organized as follows. In Section 2, we introduce the basic notations and give the definition of the objects used in later parts. Section 3 describes the connection between copulas of order statistics with fixed marginals, and symmetric copulas with fixed multidagonals. In Section 4, we determine the maximum entropy copula with fixed multidagonal. Since we obtain a symmetric copula as a result, this is also the maximum entropy symmetric copula with fixed multidagonal. In Section 5, we use the one-to-one map between the two sets of copulas established in Section 3 to give the maximum entropy copula of order statistics with fixed marginals. We finally obtain the density of the maximum entropy distribution for order statistics with fixed marginals by composing the maximum entropy copula with the marginals. Section 6 contains the detailed proofs of Theorem 4.7 and other results from Section 4. Section 7 collects the main notations of the paper to facilitate reading.

2. NOTATIONS AND DEFINITIONS

2.1. Notations in $\mathbb{R}^d$ and generalized inverse. For a Borel set $A \subset \mathbb{R}^d$, we write $|A|$ for its Lebesgue measure. For $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ and $y = (y_1, \dots, y_d) \in \mathbb{R}^d$, we write $x \leq y$ if $x_i \leq y_i$ for all $1 \leq i \leq d$. We define $\min x = \min\{x_i, 1 \leq i \leq d\}$ and $\max x = \max\{x_i, 1 \leq i \leq d\}$ for $x = (x_1, \dots, x_d) \in \mathbb{R}^d$. If J is a real-valued function defined on $\mathbb{R}$, we set $J(x) = (J(x_1), \dots, J(x_d))$. We shall consider the following subsets of $\mathbb{R}^d$:

$$S = \{(x_1, \dots, x_d) \in \mathbb{R}^d, x_1 \leq \dots \leq x_d\} \quad \text{and} \quad \Delta = S \cap I^d,$$

with $I = [0, 1]$. In what follows, usually x, y will belongs to $\mathbb{R}^d$, and s, t to $\mathbb{R}$ or I . For a set $A \subset \mathbb{R}$, we note by $A^c = \mathbb{R} \setminus A$ its complementary set.

If J is a bounded non-decreasing càd-làg function defined on $\mathbb{R}$. Its generalized inverse J^{-1} is given by $J^{-1}(t) = \inf\{s \in \mathbb{R}; J(s) \geq t\}$, for $t \in \mathbb{R}$, with the convention that $\inf \emptyset = +\infty$ and $\inf \mathbb{R} = -\infty$. We have for $s, t \in \mathbb{R}$:

$$(2) \quad J(t) \geq s \Leftrightarrow t \geq J^{-1}(s), \quad J^{-1} \circ J(t) \leq t \quad \text{and} \quad J \circ J^{-1} \circ J(t) = J(t).$$

We define the set of points where J is increasing on their left:

$$(3) \quad I_g(J) = \{t \in \mathbb{R}; u < t \Leftrightarrow J(u) < J(t)\}.$$

We have:

$$(4) \quad \mathbf{1}_{(I_g(J))^c} dJ = 0 \quad \text{a.e.},$$

$$(5) \quad J^{-1}(\mathbb{R}) \subset I_g(J) \cup \{\pm\infty\}$$

and for $s \in \mathbb{R}, t \in I_g(J)$:

$$(6) \quad J(t) \leq s \Leftrightarrow t \leq J^{-1}(s) \quad \text{and} \quad J^{-1} \circ J(t) = t.$$

Notice that if J is continuous in addition, then we have for $t \in J(\mathbb{R})$:

$$(7) \quad J \circ J^{-1}(t) = t.$$

2.2. Cdf and copula. Let $X = (X_1, \dots, X_d)$ be a random vector on $\mathbb{R}^d$. Its cumulative distribution function (cdf), denoted by F is defined by: $F(x) = \mathbb{P}(X \leq x)$, $x \in \mathbb{R}^d$. The corresponding one-dimensional marginals cdf are $(F_i, 1 \leq i \leq d)$ with $F_i(t) = \mathbb{P}(X_i \leq t)$, $t \in \mathbb{R}$. The cdf F is called a copula if X_i is uniform on $I = [0, 1]$ for all $1 \leq i \leq d$. (Notice a copula is characterized by its values on I^d only.)

We define $\mathcal{L}_d$ the set of cdf on $\mathbb{R}^d$, $\mathcal{L}_d^{1c} \subset \mathcal{L}_d$ the subset of cdf whose one-dimensional marginals cdf are continuous, $\mathcal{C} \subset \mathcal{L}_d^{1c}$ the subset of copulas. We set $\mathcal{L}_d^0$ (resp. $\mathcal{C}^0$) the subset of absolutely continuous cdf (resp. copulas) on $\mathbb{R}^d$.

Let us define for a cdf F with one-dimensional marginals $(F_i, 1 \leq i \leq d)$ the function C_F defined on I^d :

$$(8) \quad C_F(y) = F(F_1^{-1}(y_1), \dots, F_d^{-1}(y_d)), \quad y = (y_1, \dots, y_d) \in I^d.$$

If $F \in \mathcal{L}_d^{1c}$, then C_F defined by (8) is a copula thanks to (7). According to Sklar's theorem, F is then completely characterized by its one-dimensional marginals cdf $(F_i, 1 \leq i \leq d)$ and the associated copula C_F which contains all information on the dependence:

$$(9) \quad F(x) = C_F(F_1(x_1), \dots, F_d(x_d)), \quad x = (x_1, \dots, x_d) \in \mathbb{R}^d.$$

Equivalently, if $X = (X_1, \dots, X_d)$ has cdf F , then C_F is the cdf of the random vector:

$$(10) \quad (F_1(X_1), \dots, F_d(X_d)).$$

2.3. Order statistics. For $F \in \mathcal{L}_d$, we write $\mathbb{P}_F$ the distribution of a random vector $X = (X_1, \dots, X_d)$ with cdf F . A cdf $F \in \mathcal{L}_d$ is a cdf of order statistics (and we shall say that X is a vector of order statistics) if $\mathbb{P}_F(X_1 \leq X_2 \leq \dots \leq X_d) = 1$. Let us denote by $\mathcal{L}_d^{OS} \subset \mathcal{L}_d^{1c}$ the set of all cdf of order statistics with continuous one-dimensional marginals cdf. The d -tuples $(F_i, 1 \leq i \leq d)$ of marginal cdf's then verify $F_{i-1} \geq F_i$ for all $2 \leq i \leq d$. Let $\mathcal{F}_d$ be the set of d -tuples of continuous one-dimensional cdf's compatible with the marginals cdf of order statistics:

$$(11) \quad \mathcal{F}_d = \{\mathbf{F} = (F_i, 1 \leq i \leq d) \in (\mathcal{L}_1^{1c})^d; \quad \mathbf{F}_{i-1} \geq \mathbf{F}_i, \forall 2 \leq i \leq d\}.$$

For a given $\mathbf{F} = (\mathbf{F}_i, 1 \leq i \leq d)$ in $\mathcal{F}_d$, we define the set of cdf's F of order statistics with marginals cdf $\mathbf{F}$:

$$(12) \quad \mathcal{L}_d^{OS}(\mathbf{F}) = \{F \in \mathcal{L}_d^{OS}; \quad F_i = \mathbf{F}_i, 1 \leq i \leq d\}.$$

If $\mathbf{F} \in \mathcal{F}_d$, then we have $\mathcal{L}_d^{OS}(\mathbf{F}) \neq \emptyset$, since the cdf of $(\mathbf{F}_1^{-1}(U), \dots, \mathbf{F}_d^{-1}(U))$, U uniformly distributed on I , belongs to $\mathcal{L}_d^{OS}(\mathbf{F})$. We define $\mathcal{C}^{OS}(\mathbf{F})$ the set of copulas of order statistics with marginals $\mathbf{F}$:

$$(13) \quad \mathcal{C}^{OS}(\mathbf{F}) = \{C_F \in \mathcal{C}; F \in \mathcal{L}_d^{OS}(\mathbf{F})\}.$$

According to Sklar's theorem, the map $F \mapsto C_F$ is a bijection between $\mathcal{L}_d^{OS}(\mathbf{F})$ and $\mathcal{C}^{OS}(\mathbf{F})$ if $\mathbf{F} \in \mathcal{F}_d$.

2.4. Entropy. The Shannon-entropy for a cdf $F \in \mathcal{L}_d$ is given by:

$$(14) \quad H(F) = \begin{cases} -\infty & \text{if } F \in \mathcal{L}_d \setminus \mathcal{L}_d^0, \\ -\int_{\mathbb{R}^d} f \log(f) & \text{if } F \in \mathcal{L}_d^0, \end{cases}$$

with f the density of F . We will use the notation $H(X) = H(F)$ if X is a random vector with cdf F and $H(f) = H(F)$ if F has density f . The entropy of any $F \in \mathcal{L}_d^{1c}$ can be decomposed into the entropy of the one-dimensional marginals cdf $(F_i, 1 \leq i \leq d)$ and the entropy of the associated copula C_F , (see [19]):

$$(15) \quad H(F) = \sum_{i=1}^d H(F_i) + H(C_F).$$

For $\mathbf{F} = (\mathbf{F}_i, 1 \leq i \leq d) \in \mathcal{F}_d$, we define $\mathbb{J}(\mathbf{F})$ taking values in $[0, +\infty]$ by:

$$(16) \quad \mathbb{J}(\mathbf{F}) = \sum_{i=2}^d \int_{\mathbb{R}} \mathbf{F}_i(dt) |\log(\mathbf{F}_{i-1}(t) - \mathbf{F}_i(t))|.$$

Our aim is to find the cdf $F^* \in \mathcal{L}_d^{OS}(\mathbf{F})$ which maximizes the entropy H . We shall see that this is possible if and only if $\mathbb{J}(\mathbf{F})$ is finite. From an information theory point of view, this is the distribution which is the least informative among distributions of order statistics with given one-dimensional marginals cdf $\mathbf{F}$. Since the vector of marginal distribution functions $\mathbf{F}$ is fixed, thanks to (15), we notice that $H(F)$ is maximal on $\mathcal{L}_d^{OS}(\mathbf{F})$ if and only if $H(C_F)$ is maximal on $\mathcal{C}^{OS}(\mathbf{F})$. Therefore we focus on finding the copula $C^* \in \mathcal{C}^{OS}(\mathbf{F})$ which maximizes the entropy H . We will give the solution of this problem in Section 5 under some additional hypotheses on $\mathbf{F}$.

3. SYMMETRIC COPULAS WITH GIVEN ORDER STATISTICS

In this Section, we introduce an operator on the set $\mathcal{C}^{OS}(\mathbf{F})$ of copulas of order statistics with fixed marginals cdf $\mathbf{F}$. This operator assigns to a copula $C \in \mathcal{C}^{OS}(\mathbf{F})$ the copula of the exchangeable random vector associated to the order statistics with marginals cdf $\mathbf{F}$ and copula C . We show that this operator is a bijection between $\mathcal{C}^{OS}(\mathbf{F})$ and a set of symmetric copulas which can be characterized by their multidagonal, which is a generalization of the well-known diagonal section of copulas. This bijection has good properties with respect to the entropy H , giving us a problem equivalent to maximizing H on $\mathcal{C}^{OS}(\mathbf{F})$. We shall solve this problem in Section 4.

3.1. Symmetric copulas. For $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ we define $x^{OS} = (x_{(1)}, \dots, x_{(d)})$ the ordered vector (increasing order) of x , where $x_{(1)} \leq \dots \leq x_{(d)}$ and $\sum_{i=1}^d \hat{\delta}_{x_i} = \sum_{i=1}^d \hat{\delta}_{x_{(i)}}$, with $\hat{\delta}_t$ the Dirac mass at $t \in \mathbb{R}$.

Let $\mathcal{S}_d$ be the set of permutations on $\{1, \dots, d\}$. For $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ and $\pi \in \mathcal{S}_d$, we set $x_\pi = (x_{\pi(1)}, \dots, x_{\pi(d)})$. A function h defined on $\mathbb{R}^d$ is symmetric if $h(x_\pi) = h(x)$ for all $\pi \in \mathcal{S}_d$. A random vector X taking values in $\mathbb{R}^d$ is exchangeable if X_π is distributed as X for all $\pi \in \mathcal{S}_d$. In particular a random vector X taking values in $\mathbb{R}^d$ is exchangeable if and only if its cdf is symmetric. Let $\mathcal{L}_d^{sym}$ (resp. $\mathcal{C}^{sym}$) denote the subset of $\mathcal{L}_d$ (resp. $\mathcal{C}$) of symmetric cdf (resp. copulas) on $\mathbb{R}^d$.

Let $F \in \mathcal{L}_d^{1c}$ and define its symmetrization $F^{sym} \in \mathcal{L}_d^{sym}$ by:

$$(17) \quad F^{sym}(x) = \frac{1}{d!} \sum_{\pi \in \mathcal{S}_d} F(x_\pi), \quad x \in \mathbb{R}^d.$$

In particular, if X is a random vector taking values in $\mathbb{R}^d$ with cdf F and Π is a random variable independent of X , uniformly distributed on $\mathcal{S}_d$, then X_Π is exchangeable with cdf F^{sym} .

We define the following operator on the set of copulas of order statistics.

Definition 3.1. Let $\mathbf{F} \in (\mathcal{L}_1^{1c})^d$. For $C \in \mathcal{C}$ we define $S_{\mathbf{F}}(C)$ as the copula of the exchangeable random variable X_Π , where X is a random vector on $\mathbb{R}^d$ with one-dimensional marginals cdf $\mathbf{F}$ and copula C and Π is an independent random variable uniform on $\mathcal{S}_d$.

The application $S_{\mathbf{F}}$ is well-defined on $\mathcal{C}$ and takes values in $\mathcal{C}^{sym}$. In the above definition, with $\mathbf{F} = (\mathbf{F}_i, 1 \leq i \leq d)$, the one-dimensional marginals cdf of X_Π are equal to:

$$(18) \quad G = \frac{1}{d} \sum_{i=1}^d \mathbf{F}_i.$$

Since the one-dimensional marginals cdf $\mathbf{F}_i$ are continuous, we get that G is continuous and thus the cdf of X_Π belongs to $\mathcal{L}_d^{1c}$. In particular, thanks to Sklar's theorem, the copula of X_Π is indeed uniquely defined.

Combining (9), (8) and (17), we can give an explicit formula for $S_{\mathbf{F}}(C)$:

$$(19) \quad S_{\mathbf{F}}(C)(u) = \frac{1}{d!} \sum_{\pi \in \mathcal{S}_d} C(\mathbf{F}_1(G^{-1}(u_{\pi(1)})), \dots, \mathbf{F}_d(G^{-1}(u_{\pi(d)}))), \quad u \in I^d.$$

Remark 3.2. The copula $S_{\mathbf{F}}(C)$ is not equal in general to the exchangeable copula C^{sym} defined similarly to (17) by $C^{sym} = (1/d!) \sum_{\pi \in \mathcal{S}_d} C(x_\pi)$. However this is the case if the one-dimensional marginals cdf $\mathbf{F}_i$ are all equal, in which case $\mathbf{F}_i = G$ for all $1 \leq i \leq d$.

If X is a random vector on $\mathbb{R}^d$, let $X^{OS} = (X_{(1)}, \dots, X_{(d)})$ be the order statistics of X . The proof of the next Lemma is elementary.

Lemma 3.3. Let X be a random vector on $\mathbb{R}^d$ with cdf $F \in \mathcal{L}_d$ and Π a random variable independent of X , uniformly distributed on $\mathcal{S}_d$. We have:

- If $F \in \mathcal{L}_d^{OS}$, then a.s. $(X_\Pi)^{OS} = X$
- If $F \in \mathcal{L}_d^{sym}$, then $(X^{OS})_\Pi$ has the same distribution as X .

For $\mathbf{F} \in \mathcal{F}_d$, we define the set of copulas $\mathcal{C}^{sym}(\mathbf{F}) \subset \mathcal{C}^{sym}$ as the image of $\mathcal{C}^{OS}(\mathbf{F})$ by the symmetrizing operator $S_{\mathbf{F}}$:

$$(20) \quad \mathcal{C}^{sym}(\mathbf{F}) = S_{\mathbf{F}}(\mathcal{C}^{OS}(\mathbf{F})).$$

The following Lemma is one of the main result of this section.

Lemma 3.4. *Let $\mathbf{F} \in \mathcal{F}_d$. The symmetrizing operator $S_{\mathbf{F}}$ is a bijection from $\mathcal{C}^{OS}(\mathbf{F})$ onto $\mathcal{C}^{sym}(\mathbf{F})$.*

Proof. Let $C_1, C_2 \in \mathcal{C}^{OS}(\mathbf{F})$ with $S_{\mathbf{F}}(C_1) = S_{\mathbf{F}}(C_2)$. Let X and Y be random vectors with one-dimensional marginals cdf $\mathbf{F}$ and copula C_1, C_2 respectively. Since $C_1, C_2 \in \mathcal{C}^{OS}(\mathbf{F})$, we get that X and Y are order statistics. Notice X_{Π} and Y_{Π} have the same one-dimensional marginals according to (18) and same copula given by $S_{\mathbf{F}}(C_1) = S_{\mathbf{F}}(C_2)$. Therefore X_{Π} and Y_{Π} have the same distribution. Thus, their corresponding order statistics $(X_{\Pi})^{OS}$ and $(Y_{\Pi})^{OS}$ have the same distribution. By Lemma 3.3 we get that X and Y have the same distribution as well, which implies $C_1 = C_2$. $\square$

Remark 3.5. We have in general $\mathcal{C}^{sym}(\mathbf{F}) \neq \mathcal{C}^{OS}(\mathbf{F}) \cap \mathcal{C}^{sym}$. One exception being when the marginals cdf's $\mathbf{F}_i$ are all equal. In this case, both sides reduce to one copula which is the Fréchet-Hoeffding upper bound copula: $C^+(u) = \min u$, $u \in I^d$.

3.2. Multidiagonals and characterization of $\mathcal{C}^{sym}(\mathbf{F})$. Let $C \in \mathcal{C}$ be a copula and U a random vector with cdf C . The map $t \mapsto C(t, \dots, t)$ for $t \in I$, which is called the diagonal section of C , is the cdf of $\max U$. We shall consider a generalization of the diagonal section of C in the next Definition.

Definition 3.6. *Let $C \in \mathcal{C}$ be a copula on $\mathbb{R}^d$ and U a random vector with cdf C . The multidagonal of the copula C , $\delta_C = (\delta_{(i)}, 1 \leq i \leq d)$, is the d -tuple of the one-dimensional marginals cdf of $U^{OS} = (U_{(1)}, \dots, U_{(d)})$ the order statistics of U : for $1 \leq i \leq d$*

$$\delta_{(i)}(t) = \mathbb{P}(U_{(i)} \leq t), \quad t \in I.$$

We denote by $\mathcal{D} = \{\delta_C; C \in \mathcal{C}\}$ the set of multidagonals. Notice that $\mathcal{D} \subset \mathcal{F}_d$, see Remark 3.7. For $\delta \in \mathcal{D}$ a multidagonal, we define $\mathcal{C}_{\delta} = \{C; \delta_C = \delta\}$ the set of copulas with multidagonal δ .

A characterization of the set $\mathcal{D}$ is given by Theorem 1 of [12]: a vector of functions $\delta = (\delta_{(1)}, \dots, \delta_{(d)})$ belongs to $\mathcal{D}$ if and only if $\delta_{(i)} \in \mathcal{L}_1$ and the following conditions hold:

$$(21) \quad \delta_{(i)} \geq \delta_{(i+1)}, \quad 1 \leq i \leq d-1,$$

$$(22) \quad \sum_{i=1}^d \delta_{(i)}(s) = ds, \quad 0 \leq s \leq 1.$$

Remark 3.7. The condition (22) implies that $\delta_{(i)} \in \mathcal{L}_1^{1c}$, $1 \leq i \leq d$, moreover they are d -Lipschitz. Also, it is enough to know $d-1$ functions from $\delta_{(i)}$, $1 \leq i \leq d$, the remaining one is implicitly defined by (22). Condition (21) along with the continuity of $\delta_{(i)}$ implies that any multidagonal δ_C is compatible with the continuous marginal distributions of an order statistics, therefore $\mathcal{D} \subset \mathcal{F}_d$.

Remark 3.8. Since $\delta_{(i)}$, $1 \leq i \leq d$ are non-decreasing and d -Lipschitz, we have for almost every $t \in I$: $0 \leq (\delta_{(i)})'(t) \leq d$ and thus $|(\delta_{(i)})'(t) \log((\delta_{(i)})'(t))| \leq d \log(d)$ for $d \geq 2$. We deduce that for $d \geq 2$:

$$(23) \quad |H(\delta_{(i)})| \leq d \log(d).$$

Remark 3.9. Let $C \in \mathcal{C}^{sym}$ be a symmetric copula on $\mathbb{R}^d$ and U a random vector with cdf C . We check that the multidagonal $\delta_C = (\delta_{(i)}, 1 \leq i \leq d)$ can be expressed in terms of the diagonal sections $(C_{\{i\}}, 1 \leq i \leq d)$ where for $1 \leq i \leq d$:

$$C_{\{i\}}(t) = \mathbb{P} \left(\max_{1 \leq k \leq i} U_k \leq t \right) = C(\underbrace{t, \dots, t}_i \text{ terms}, \underbrace{1, \dots, 1}_{d-i \text{ terms}}), \quad t \in I.$$

According to 2.8 of [12], we have for $1 \leq i \leq d$:

$$\delta_{(i)}(t) = \sum_{j=i}^d (-1)^{j-i} \binom{j-1}{i-1} \binom{d}{j} C_{\{j\}}(t), \quad t \in I.$$

Conversely, we can express the functions $(C_{\{i\}}, 1 \leq i \leq d)$ with δ_C . For $1 \leq i \leq d$ and $t \in I$, we have:

$$\begin{aligned} C_{\{i\}}(t) &= \mathbb{P} \left(\max_{1 \leq k \leq i} U_k \leq t \right) \\ &= \sum_{j=i}^d \mathbb{P} \left(U_{(j)} \leq t \mid \max_{1 \leq k \leq i} U_k = U_{(j)} \right) \mathbb{P} \left(\max_{1 \leq k \leq i} U_k = U_{(j)} \right) \\ &= \sum_{j=i}^d \frac{\binom{j-1}{i-1}}{\binom{d}{i}} \delta_{(j)}(t), \end{aligned}$$

where we used the definition of $\delta_{(i)}$ and the exchangeability of U for the third equality.

The next technical Lemma will be used in forthcoming proofs. Recall that J^{-1} denotes the generalized inverse of a non-decreasing function J , see Section 2.1 for its definition and properties, in particular, $J^{-1} \circ J(t) \leq t$ for $t \in \mathbb{R}$. Recall also that for $x = (x_1, \dots, x_d) \in \mathbb{R}^d$, we write $G(x) = (G(x_1), \dots, G(x_d))$.

Lemma 3.10. *Let $X = (X_1, \dots, X_d)$ be a random vector on $\mathbb{R}^d$ with one-dimensional marginals cdf $(F_i, 1 \leq i \leq d) \in (\mathcal{L}_1)^d$. Set $G = \sum_{i=1}^d F_i/d$. We have for $1 \leq i \leq d$:*

$$(24) \quad \mathbb{P}(X_i \leq G^{-1} \circ G(t)) = \mathbb{P}(X_i \leq t), \quad t \in \mathbb{R}, \quad \text{that is} \quad F_i \circ G^{-1} \circ G = F_i.$$

We also have for $x \in \mathbb{R}^d$:

$$(25) \quad \mathbb{P}(G(X) \leq x) = \mathbb{P}(X \leq G^{-1}(x)).$$

Proof. Since G is the average of the non-decreasing functions F_i , if $G(s) = G(s')$ for some $s, s' \in \mathbb{R}$, then we have $F_i(s) = F_i(s')$ for every $1 \leq i \leq d$. Thanks to (2), we have $G \circ G^{-1} \circ G(t) = G(t)$ and thus $F_i \circ G^{-1} \circ G(t) = F_i(t)$. This gives (24).

Recall definition (3) for $I_g(J)$ the set of points where the function J is increasing on their left. Since G is the average of the non-decreasing functions F_i , we deduce that $I_g(G) = \bigcup_{1 \leq i \leq d} I_g(F_i)$. Notice that a.s. X_i belongs to $I_g(F_i)$. Thanks to (6), we get that a.s. $\{G(X) \leq x\} = \{X \leq G^{-1}(x)\}$. This gives (25). $\square$

We will also require the following Lemma.

Lemma 3.11. *Let $X = (X_1, \dots, X_d)$ be a random vector on $\mathbb{R}^d$ with one-dimensional marginals cdf $(F_i, 1 \leq i \leq d) \in (\mathcal{L}_1)^d$. Set $G = \sum_{i=1}^d F_i/d$. We have for $1 \leq i \leq d$:*

$$(26) \quad (F_i \circ G^{-1})^{-1} = G \circ F_i^{-1}.$$

Proof. Recall Definition (3) for $I_g(J)$ the set of points where the function J is increasing on their left. Let $1 \leq i \leq d$. Thanks to (5), we have $F_i^{-1}(\mathbb{R}) \subset I_g(F_i) \cup \{\pm\infty\}$. Since G is the average of the non-decreasing functions F_i , we deduce that $I_g(G) = \bigcup_{1 \leq i \leq d} I_g(F_i)$. Thus we get:

$$(27) \quad F_i^{-1}(\mathbb{R}) \subset I_g(G) \cup \{\pm\infty\},$$

for all $1 \leq i \leq d$. The function $F_i \circ G^{-1}$ is also bounded, non-decreasing and càd-làg therefore we have for $t, s, \in \mathbb{R}$:

$$t \geq (F_i \circ G^{-1})^{-1}(s) \iff F_i \circ G^{-1}(t) \geq s \iff G^{-1}(t) \geq F_i^{-1}(s) \iff t \geq G \circ F_i^{-1}(s),$$

where we used the equivalence of (2) for the first and second equivalence, (27) and the equivalence of (6) for the last. This gives that $(F_i \circ G^{-1})^{-1} = G \circ F_i^{-1}$. $\square$

In the following Lemma, we show that for $\mathbf{F} \in \mathcal{F}_d$, all copulas in $\mathcal{C}^{sym}(\mathbf{F})$ share the same multidagonal denoted by $\delta^{\mathbf{F}}$.

Lemma 3.12. *Let $\mathbf{F} = (\mathbf{F}_i, 1 \leq i \leq d) \in \mathcal{F}_d$. Let $C \in \mathcal{C}^{OS}(\mathbf{F})$ and U be a random vector with cdf $S_{\mathbf{F}}(C)$. Let $\delta^{\mathbf{F}} = (\delta_{(i)}, 1 \leq i \leq d)$ be the multidagonal of $S_{\mathbf{F}}(C)$, that is the one-dimensional marginals cdf of U^{OS} , the order statistics of U . We have that $\delta^{\mathbf{F}}$ does not depend on C and for $1 \leq i \leq d$:*

$$(28) \quad \delta_{(i)} = \mathbf{F}_i \circ G^{-1} \quad \text{and} \quad \delta_{(i)}^{-1} = G \circ \mathbf{F}_i^{-1},$$

with G given by (18). Furthermore, C is the unique copula of U^{OS} .

With obvious notation, we might simply write $\delta^{\mathbf{F}} = \mathbf{F} \circ G^{-1}$, with G given by (18).

Proof. Let X be a random vector of order statistics with marginals $\mathbf{F} \in \mathcal{F}_d$ and copula C . Then $S_{\mathbf{F}}(C)$ is the copula of the exchangeable random vector X_{Π} , where Π is uniform on $\mathcal{S}_d$ and independent of X . We have already seen in (18) that the one-dimensional marginals of X_{Π} have the same distribution given by $G \in \mathcal{L}_1^{1c}$. Thanks to (10), we deduce that the random vector U , with cdf $S_{\mathbf{F}}(C)$, has the same distribution as $G(X_{\Pi})$. Since G is non-decreasing, this implies that the order statistics of U , U^{OS} , has the same distribution as $G((X_{\Pi})^{OS})$ that is as $G(X)$, thanks to Lemma 3.3. Then use (25) to get for $x \in \mathbb{R}^d$:

$$(29) \quad \mathbb{P}(U^{OS} \leq x) = \mathbb{P}(G(X) \leq x) = \mathbb{P}(X \leq G^{-1}(x)).$$

This gives the first part of the Lemma as the multidagonal of U is the one-dimensional marginals cdf of its order statistics. The second equation in (28) is due to Lemma 3.11. The fact that C is the copula of U^{OS} and its uniqueness are due to (29) and the continuity of $\delta_{(i)}$, see Remark 3.7. $\square$

According to the next Proposition, (20) and Lemma 3.4, we get the main result of this Section: for any $\mathbf{F} \in \mathcal{F}_d$, the symmetrizing operator $S_{\mathbf{F}}$ is a bijection between $\mathcal{C}^{OS}(\mathbf{F})$ and $\mathcal{C}_{\delta^{\mathbf{F}}} \cap \mathcal{C}^{sym}$.

Proposition 3.13. *Let $\mathbf{F} \in \mathcal{F}_d$. We have $\mathcal{C}^{sym}(\mathbf{F}) = \mathcal{C}_{\delta^{\mathbf{F}}} \cap \mathcal{C}^{sym}$.*

Proof. By Lemma 3.12, we have $\mathcal{C}^{sym}(\mathbf{F}) \subset \mathcal{C}_{\delta^{\mathbf{F}}} \cap \mathcal{C}^{sym}$.

Let $C \in \mathcal{C}_{\delta^{\mathbf{F}}} \cap \mathcal{C}^{sym}$ and U be a random vector with cdf C . Let G be given by (18). Notice that $X = G^{-1}(U)$ is an exchangeable random vector with marginals G and copula C . Thanks to Lemma 3.3, the proof will be complete as soon as we prove that the one-dimensional marginals cdf of $X^{OS} = (X_{(1)}, \dots, X_{(d)})$, the order statistics of X , is given by

F. Notice $X^{OS} = G^{-1}(U^{OS})$, with U^{OS} the order statistics of U whose one-dimensional marginals cdf are given by $\delta^{\mathbf{F}}$. We have for $1 \leq i \leq d$ and $t \in \mathbb{R}$:

$$\mathbb{P}(X_{(i)} \leq t) = \mathbb{P}(G^{-1}(U_{(i)}) \leq t) = \mathbb{P}(U_{(i)} \leq G(t)) = \mathbf{F}_i \circ G^{-1} \circ G(t) = \mathbf{F}_i(t),$$

where we used (2) for the second equality, (28) for the third, and (24) for the last. This finishes the proof. $\square$

We end this Section by an ancillary result we shall use later.

Lemma 3.14. *Let $\mathbf{F} \in \mathcal{F}_d$. We have $\mathbb{J}(\mathbf{F}) = \mathbb{J}(\delta^{\mathbf{F}})$.*

Proof. Let $\mathbf{F} = (\mathbf{F}_i, 1 \leq i \leq d)$. We get, using (28) and the change of variable $s = G^{-1}(t)$ that:

$$\int_I \delta'_{(i)}(t) |\log(\delta_{(i-1)}(t) - \delta_{(i)}(t))| dt = \int_{G^{-1}((0,1))} \mathbf{F}_i(ds) |\log(\mathbf{F}_{i-1}(s) - \mathbf{F}_i(s))|.$$

Since $d\mathbf{F}_i = 0$ outside $G^{-1}((0,1))$ (as G is increasing as soon as $\mathbf{F}_i$ is increasing), we get the above last integration is also over $\mathbb{R}$. We deduce that:

$$\mathbb{J}(\delta^{\mathbf{F}}) = \sum_{i=2}^d \int_{\mathbb{R}} \mathbf{F}_i(ds) |\log(\mathbf{F}_{i-1}(s) - \mathbf{F}_i(s))| = \mathbb{J}(\mathbf{F}).$$

$\square$

3.3. Density and entropy of copulas in $\mathcal{C}^{sym}(\mathbf{F})$. We prove in this Section that $S_{\mathbf{F}}$ preserves the absolute continuity on $\mathcal{C}^{OS}(\mathbf{F})$ for $\mathbf{F} \in \mathcal{F}_d$ and the entropy up to a constant. Let us introduce some notation. For marginals $\mathbf{F} \in \mathcal{F}_d$, let

$$(30) \quad \Psi_i^{\mathbf{F}} = \{s \in \mathbb{R}, \mathbf{F}_{i-1}(s) > \mathbf{F}_i(s)\} \quad \text{for } 2 \leq i \leq d.$$

The complementary set $(\Psi_i^{\mathbf{F}})^c$ is the collection of the points where $\mathbf{F}_{i-1} = \mathbf{F}_i$. We define $\Sigma^{\mathbf{F}} \subset I$ as:

$$(31) \quad \Sigma^{\mathbf{F}} = \bigcup_{i=2}^d \mathbf{F}_i((\Psi_i^{\mathbf{F}})^c).$$

By Remark 3.7 we have $\mathcal{D} \subset \mathcal{F}_d$, then the definitions (30) and (31) apply for all $\delta \in \mathcal{D}$. In particular, for $\delta = (\delta_{(1)}, \dots, \delta_{(d)}) \in \mathcal{D}$ the sets Ψ_i^{δ} , $2 \leq i \leq d$ are open subsets of I , therefore $(\Psi_i^{\delta})^c \cap I$ is a compact subset. This and the continuity of $\delta_{(i)}$ imply that $\delta_{(i)}((\Psi_i^{\delta})^c) = \delta_{(i)}((\Psi_i^{\delta})^c \cap I)$ is also compact, hence Σ^{δ} is compact. Notice that $\{0, 1\} \subset \Sigma^{\delta}$ always holds. We define $\mathcal{C}_{\delta}^0 = \mathcal{C}_{\delta} \cap \mathcal{C}^0$ the subset of absolutely continuous copulas with multidagonal δ and the subset $\mathcal{D}^0 = \{\delta \in \mathcal{D}, \mathcal{C}_{\delta}^0 \neq \emptyset\}$ of multidagonals of absolutely continuous copulas. According to Theorem 2 of [12], the multidagonal δ belongs to $\mathcal{D}^0$ if and only if it belongs to $\mathcal{D}$ and the Lebesgue measure of Σ^{δ} is zero: $|\Sigma^{\delta}| = 0$.

Lemma 3.15. *Let $\delta \in \mathcal{D}$. We have $\delta \in \mathcal{D}^0$ if and only if for all $2 \leq i \leq d$, a.e.:*

$$(32) \quad \delta'_{(i-1)} \mathbf{1}_{(\Psi_i^{\delta})^c} = \delta'_{(i)} \mathbf{1}_{(\Psi_i^{\delta})^c} = 0.$$

Furthermore, we have that $\mathbb{J}(\delta) < +\infty$ implies $\delta \in \mathcal{D}^0$.

Proof. Let J be a function defined on I , Lipschitz and non-decreasing. Let A be a Borel subset of I . We have:

$$|J(A)| = \int \mathbf{1}_{J(A)}(t) dt = \int_0^1 \mathbf{1}_{\{s \in J^{-1} \circ J(A)\}} J'(s) ds = \int_0^1 \mathbf{1}_A(s) J'(s) ds,$$

where we used (4) and (6) for the last equality. This gives that $|J(A)| = 0$ if and only if a.e. $J' \mathbf{1}_A = 0$. Then use that $\delta \in \mathcal{D}^0$ if and only if $|\delta_{(i)}((\Psi_i^\delta)^c)| = 0$ for all $1 \leq i \leq d$ and that $\delta_{(i-1)}((\Psi_i^\delta)^c) = \delta_{(i)}((\Psi_i^\delta)^c)$ to conclude that $\delta \in \mathcal{D}^0$ if and only if (32) holds for all $2 \leq i \leq d$. The last part of the Lemma is clear. $\square$

Definition 3.16. Let $\mathcal{F}_d^0 \subset \mathcal{F}_d$ be the subset of marginals $\mathbf{F}$ such that there exists an absolutely continuous cdf of order statistics with marginals $\mathbf{F}$, that is $\mathcal{L}_d^{OS}(\mathbf{F}) \cap \mathcal{L}_d^0 \neq \emptyset$.

In particular, we have $\mathcal{D}^0 \subset \mathcal{F}_d^0$. The next Lemma gives a characterization of the set $\mathcal{F}_d^0$.

Lemma 3.17. Let $\mathbf{F} \in \mathcal{F}_d$. Then $\mathbf{F} \in \mathcal{F}_d^0$ if and only if $\mathbf{F}_i \in \mathcal{L}_1^0$ for $1 \leq i \leq d$ and $|\Sigma^{\mathbf{F}}| = 0$. Furthermore, we have that $\mathbf{F}_i \in \mathcal{L}_1^0$ for $1 \leq i \leq d$ and $\mathbb{J}(\mathbf{F}) < +\infty$ imply $\mathbf{F} \in \mathcal{F}_d^0$.

Proof. Let $F \in \mathcal{L}_d^{OS}(\mathbf{F})$. We know that $F \in \mathcal{L}_d^0$ if and only if $\mathbf{F}_i \in \mathcal{L}_1^0$ for $1 \leq i \leq d$ and $C_F \in \mathcal{C}^0$, the subset of absolutely continuous copulas (see for example [11]). Therefore $\mathbf{F} \in \mathcal{F}_d^0$ if and only if $\mathbf{F}_i \in \mathcal{L}_1^0$ for $1 \leq i \leq d$ and $\mathcal{C}^{OS}(\mathbf{F}) \cap \mathcal{C}^0 \neq \emptyset$. Recall that $\delta^{\mathbf{F}}$ is defined by (28). We first show that

$$(33) \quad \mathcal{C}^{OS}(\mathbf{F}) \cap \mathcal{C}^0 \neq \emptyset \text{ if and only if } \mathcal{C}_{\delta^{\mathbf{F}}}^0 \cap \mathcal{C}^{sym} \neq \emptyset.$$

Let $C \in \mathcal{C}^{OS}(\mathbf{F}) \cap \mathcal{C}^0$. Then Lemma 3.12 ensures that $S_{\mathbf{F}}(C) \in \mathcal{C}_{\delta^{\mathbf{F}}} \cap \mathcal{C}^{sym}$. The absolute continuity of $S_{\mathbf{F}}(C)$ is a direct consequence of (19), (28) and Remark 3.7 which ensures that $\delta_{(i)}^{\mathbf{F}}, 1 \leq i \leq d$ are d -Lipschitz, therefore their derivatives exist a.e. on I . This ensures that $\mathcal{C}_{\delta^{\mathbf{F}}}^0 \cap \mathcal{C}^{sym} \neq \emptyset$.

Conversely, let $C \in \mathcal{C}_{\delta^{\mathbf{F}}}^0 \cap \mathcal{C}^{sym}$. Let U be a random vector with cdf C . Then its order statistics U^{OS} is also absolutely continuous. Therefore the copula of U^{OS} , which is $S_{\mathbf{F}}^{-1}(C)$ by Lemma 3.12, is also absolutely continuous. This proves thanks to Proposition 3.13 and Lemma 3.4 that $S_{\mathbf{F}}^{-1}(C) \in \mathcal{C}^{OS}(\mathbf{F}) \cap \mathcal{C}^0$. This gives (33).

Notice that $\mathcal{C}_{\delta^{\mathbf{F}}}^0 \cap \mathcal{C}^{sym} \neq \emptyset$ is equivalent to $\mathcal{C}_{\delta^{\mathbf{F}}}^0 \neq \emptyset$, since for any $C \in \mathcal{C}_{\delta^{\mathbf{F}}}^0$ we have that C^{sym} defined by (17) belongs to $\mathcal{C}_{\delta^{\mathbf{F}}}^0 \cap \mathcal{C}^{sym}$. By Theorem 2 of [12], $\mathcal{C}_{\delta^{\mathbf{F}}}^0 \neq \emptyset$ if and only if $\Sigma^{\delta^{\mathbf{F}}}$ has zero Lebesgue measure. The proof is then complete as one can easily verify using (28) that $\Sigma^{\delta^{\mathbf{F}}} = \Sigma^{\mathbf{F}}$ and thanks to Lemma 3.14. $\square$

From now on we consider $\mathbf{F} \in \mathcal{F}_d^0$. We give an auxiliary lemma on the support of the copulas in $\mathcal{C}^{OS}(\mathbf{F}) \cap \mathcal{C}^0$.

Lemma 3.18. Let $\mathbf{F} = (\mathbf{F}_i, 1 \leq i \leq d) \in \mathcal{F}_d^0$ and $C \in \mathcal{C}^{OS}(\mathbf{F}) \cap \mathcal{C}^0$. Then the density of C vanishes a.e. on $I^d \setminus T^{\mathbf{F}}$ with:

$$(34) \quad T^{\mathbf{F}} = \{u = (u_1, \dots, u_d) \in I^d; \mathbf{F}_1^{-1}(u_1) \leq \dots \leq \mathbf{F}_d^{-1}(u_d)\}.$$

Proof. Let $X = (X_1, \dots, X_d)$ be a random vector of order statistics with one-dimensional marginals cdf $\mathbf{F}$ and copula $C \in \mathcal{C}^0$. Let $U = (U_1, \dots, U_d)$ be a random vector with cdf C . Then it is distributed as $(\mathbf{F}_1(X_1), \dots, \mathbf{F}_d(X_d))$, see (10). We get $\mathbb{P}(U \in T^{\mathbf{F}}) = 1$, since X is a vector of order statistics and $X_i \in I_g(\mathbf{F}_i)$ a.s. for $1 \leq i \leq d$. This gives the result. $\square$

Now we establish the connection between the sets $\mathcal{C}^{OS}(\mathbf{F}) \cap \mathcal{C}^0$ and $\mathcal{C}^{sym}(\mathbf{F}) \cap \mathcal{C}^0$.

Lemma 3.19. *Let $\mathbf{F} \in \mathcal{F}_d^0$. The symmetrizing operator $S_{\mathbf{F}}$ is a bijection from $\mathcal{C}^{OS}(\mathbf{F}) \cap \mathcal{C}^0$ onto $\mathcal{C}^{sym}(\mathbf{F}) \cap \mathcal{C}^0$. Moreover, if $C \in \mathcal{C}^{OS}(\mathbf{F}) \cap \mathcal{C}^0$, with density function c , then the density function $s_{\mathbf{F}}(C)$ of $S_{\mathbf{F}}(C)$ is given by, for a.e. $u = (u_1, \dots, u_d) \in I^d$:*

$$(35) \quad s_{\mathbf{F}}(C)(u) = \frac{1}{d!} c(\delta_{(1)}(u_{(1)}), \dots, \delta_{(d)}(u_{(d)})) \prod_{i=1}^d \delta'_{(i)}(u_{(i)}).$$

Let $T^{\mathbf{F}}$ be given by (34). If $C \in \mathcal{C}^{sym}(\mathbf{F}) \cap \mathcal{C}_0$ with density c , then the density $s_{\mathbf{F}}^{-1}(C)$ of $S_{\mathbf{F}}^{-1}(C)$ is given by, for a.e. $u = (u_1, \dots, u_d) \in I^d$:

$$(36) \quad s_{\mathbf{F}}^{-1}(C)(u) = d! \frac{c(\delta_{(1)}^{-1}(u_1), \dots, \delta_{(d)}^{-1}(u_d))}{\prod_{i=1}^d \delta'_{(i)} \circ \delta_{(i)}^{-1}(u_i)} \mathbf{1}_{T^{\mathbf{F}}}(u) \mathbf{1}_{\{\prod_{i=1}^d \delta'_{(i)} \circ \delta_{(i)}^{-1}(u_i) > 0\}}.$$

Proof. By Proposition 3.13, we deduce that $\mathcal{C}^{sym}(\mathbf{F}) \cap \mathcal{C}^0 = \mathcal{C}_{\delta_{\mathbf{F}}}^0 \cap \mathcal{C}^{sym}$. Lemma 3.4 and the proof of Lemma 3.17 ensures that $S_{\mathbf{F}}$ is a bijection between $\mathcal{C}^{OS}(\mathbf{F}) \cap \mathcal{C}^0$ and $\mathcal{C}^{sym}(\mathbf{F}) \cap \mathcal{C}^0$. The explicit formula (35) can be obtained by taking the mixed derivative of the right hand side of (19). By Lemma 3.18, all the terms in the sum disappear except the one on the right hand side of (35).

To obtain (36), let $C \in \mathcal{C}^{sym}(\mathbf{F}) \cap \mathcal{C}^0$ with density c , and U be a random vector with cdf C . The order statistics U^{OS} derived from U is also absolutely continuous with cumulative distribution function K , and density function k given by:

$$k(u) = d! c(u) \mathbf{1}_{\Delta}(u), \quad u \in I^d.$$

By Lemma 3.12, $S_{\mathbf{F}}^{-1}(C)$ is the copula of U^{OS} . From (8), we have for $u = (u_1, \dots, u_d) \in I^d$:

$$(37) \quad S_{\mathbf{F}}^{-1}(C)(u) = K(\delta_{(1)}^{-1}(u_1), \dots, \delta_{(d)}^{-1}(u_d)).$$

According to (6), we deduce that $G^{-1} \circ G \circ \mathbf{F}_i^{-1} = \mathbf{F}_i^{-1}$ on $(0, 1)$. This implies that for $s, t \in (0, 1)$, $1 \leq i < j \leq d$:

$$\begin{aligned} \delta_{(i)}^{-1}(s) \leq \delta_{(j)}^{-1}(t) &\Leftrightarrow G \circ \mathbf{F}_i^{-1}(s) \leq G \circ \mathbf{F}_j^{-1}(t) \\ &\Rightarrow G^{-1} \circ G \circ \mathbf{F}_i^{-1}(s) \leq G^{-1} \circ G \circ \mathbf{F}_j^{-1}(t) \\ &\Leftrightarrow \mathbf{F}_i^{-1}(s) \leq \mathbf{F}_j^{-1}(t) \\ &\Rightarrow G \circ \mathbf{F}_i^{-1}(s) \leq G \circ \mathbf{F}_j^{-1}(t), \end{aligned}$$

where we used (28) for the first equivalence, that G^{-1} is non-decreasing for the first implication and G is non-decreasing for the second. Thus, we have that for $s, t \in (0, 1)$, that the two conditions $\delta_{(i)}^{-1}(s) \leq \delta_{(j)}^{-1}(t)$ and $\mathbf{F}_i^{-1}(s) \leq \mathbf{F}_j^{-1}(t)$ are equivalent. Thus we deduce that the two sets

$$\left\{ (u_1, \dots, u_d) \in I^d; \delta_{(1)}^{-1}(u_1) \leq \dots \leq \delta_{(d)}^{-1}(u_d) \right\}$$

and $T^{\mathbf{F}}$ are equal up to a set of zero Lebesgue measure. Then we deduce (36) from (37). $\square$

We give a general result on the entropy of an exchangeable random vector and the entropy of its order statistics.

Lemma 3.20. *Let X be a random vector on $\mathbb{R}^d$, X^{OS} the corresponding order statistics and Π an independent uniform random variable on $\mathcal{S}_d$. Then we have:*

$$H((X^{OS})_{\Pi}) = \log(d!) + H(X^{OS}).$$

Proof. Let F be the cdf of X^{OS} . If $F \notin \mathcal{L}_d^0$, then the cdf F^{sym} of $(X^{OS})_\Pi$ given by (17) verifies also $F^{sym} \notin \mathcal{L}_d^0$, therefore $H((X^{OS})_\Pi) = H(X^{OS}) + \log(d!) = -\infty$. If $F \in \mathcal{L}_d^0$ with density function f , then the density function f^{sym} of F^{sym} is given by, for $x \in \mathbb{R}^d$:

$$f^{sym}(x) = \frac{1}{d!} f(x^{OS}),$$

where x^{OS} is the ordered vector of x . Therefore, using that $f(x) = 0$ if $x \neq x^{OS}$, we have:

$$\begin{aligned} H((X^{OS})_\Pi) &= - \int_{\mathbb{R}^d} f^{sym} \log(f^{sym}) \\ &= \log(d!) - \frac{1}{d!} \int_{\mathbb{R}^d} f(x^{OS}) \log(f(x^{OS})) dx \\ &= \log(d!) - \int_{\mathbb{R}^d} f(x) \log(f(x)) dx \\ &= \log(d!) + H(X^{OS}). \end{aligned}$$

□

Now we are ready to give the connection between the entropy of C and $S_{\mathbf{F}}(C)$ for $C \in \mathcal{C}^{OS}(\mathbf{F})$, which is the main result of this Section. Recall the definition of $\delta^{\mathbf{F}} = (\delta_{(i)}^{\mathbf{F}}, 1 \leq i \leq d)$ given in Lemma 3.12 and thanks to Remark 3.8, $H(\delta_{(i)}^{\mathbf{F}})$ is finite for all $1 \leq i \leq d$.

Proposition 3.21. *Let $\mathbf{F} \in \mathcal{F}_d$ and $C \in \mathcal{C}^{OS}(\mathbf{F})$. Then we have:*

$$(38) \quad H(S_{\mathbf{F}}(C)) = \log(d!) + H(C) + \sum_{i=1}^d H(\delta_{(i)}^{\mathbf{F}}).$$

Proof. Let U be an exchangeable random vector with cdf $S_{\mathbf{F}}(C)$, and U^{OS} its order statistics. According to Lemma 3.12, U^{OS} has one-dimensional marginals cdf $\delta^{\mathbf{F}} = (\delta_{(i)}^{\mathbf{F}}, 1 \leq i \leq d)$ and copula C . Therefore, using (15), we get:

$$H(U^{OS}) = H(C) + \sum_{i=1}^d H(\delta_{(i)}^{\mathbf{F}}).$$

On the other hand, since $S_{\mathbf{F}}(C)$ is symmetric, Lemma 3.3 ensures that $(U^{OS})_\Pi$ has the same distribution as U . Therefore Lemma 3.20 gives:

$$H(S_{\mathbf{F}}(C)) = H(U) = H((U^{OS})_\Pi) = H(U^{OS}) + \log(d!) = H(C) + \sum_{i=1}^d H(\delta_{(i)}^{\mathbf{F}}) + \log(d!).$$

□

4. MAXIMUM ENTROPY COPULA WITH GIVEN MULTIDIAGONALS

This section is a generalization of [6], where the maximum entropy copula with given diagonal section (i.e. given distribution for the maximum of its marginals) is studied.

Recall that multidagonals of copulas on $\mathbb{R}^d$ are given by Definition 3.6. We recall some further notation: $\mathcal{D}$ denotes the set of multidagonals; for $\delta \in \mathcal{D}$, $\mathcal{C}_\delta$ denotes the subset of copulas with multidagonal δ ; $\mathcal{C}^0$ denotes the subset copulas which are absolutely continuous, and $\mathcal{C}_\delta^0 = \mathcal{C}_\delta \cap \mathcal{C}^0$. The set $\mathcal{D}^0 \subset \mathcal{D}$ contains all diagonals for which $\mathcal{C}_\delta^0 \neq \emptyset$.

We give an explicit formula for C^* such that $H(C^*) = \max_{C \in \mathcal{C}_\delta} H(C)$, with H the entropy, see definition (14). Notice that the maximum can be taken over $\mathcal{C}_\delta^0$, since the entropy is minus infinity otherwise. When $d = 2$, the problem was solved in [6].

Let $\delta = (\delta_{(i)}, 1 \leq i \leq d) \in \mathcal{D}$ be a multidagonal. Since $\delta_{(i)}, 1 \leq i \leq d$ are d -Lipschitz, the entropy of $H(\delta_{(i)})$ is well defined and finite, see Remark 3.8 and $\mathbb{J}(\delta)$ given by (16) is also well defined and belongs to $[0, +\infty]$.

The next two lemmas provides sets on which the density of a copula with given multidagonal is zero. For $\delta \in \mathcal{D}$, let:

$$(39) \quad Z_\delta = \{u \in I^d; \text{ there exists } 1 \leq i \leq d \text{ such that } \delta'_{(i)}(u_{(i)}) = 0\}.$$

Lemma 4.1. *Let $\delta \in \mathcal{D}^0$. Then for all copulas $C \in \mathcal{C}_\delta^0$ with density c , we have $c \mathbf{1}_{Z_\delta} = 0$ a.e. that is $c(u) \mathbf{1}_{Z_\delta}(u) = 0$ for a.e. $u \in I^d$.*

Proof. By definition of $\delta_{(i)}$, we have for all $r \in I$:

$$\int_{I^d} c(u) \mathbf{1}_{\{u_{(i)} \leq r\}} du = \delta_{(i)}(r) = \int_0^r \delta'_{(i)}(s) ds.$$

This implies, by the monotone class theorem, that for all measurable subset K of I , we have:

$$\int_{I^d} c(u) \mathbf{1}_K(u_{(i)}) du = \int_K \delta'_{(i)}(s) ds.$$

Since $c \geq 0$ a.e., we deduce that a.e. $c(u) \mathbf{1}_{\{\delta'_{(i)}(u_{(i)})=0\}} = 0$ and thus a.e. $c \mathbf{1}_{Z_\delta} = 0$. $\square$

Recall the definition of Ψ_i^δ given by (30) for $2 \leq i \leq d$. We also define $\Psi_1^\delta = (0, d_1)$ with $d_1 = \inf\{s \in I; \delta_{(1)}(s) = 1\}$ and $\Psi_{d+1}^\delta = (g_{d+1}, 1)$ with $g_{d+1} = \sup\{s \in I; \delta_{(d)}(s) = 0\}$. Since Ψ_i^δ are open subsets of I , there exists at most countably many disjoint intervals $\{(g_i^{(j)}, d_i^{(j)})\}$, $j \in J_i$ such that

$$(40) \quad \Psi_i^\delta = \bigcup_{j \in J_i} (g_i^{(j)}, d_i^{(j)}).$$

We note by $m_i^{(j)} = (g_i^{(j)} + d_i^{(j)})/2$ the midpoint of these intervals for $2 \leq i \leq d+1$. In particular $m_{d+1} = (1 + g_{d+1})/2$. We also define $m_1 = 0$. For $\delta \in \mathcal{D}$, let:

$$(41) \quad L_\delta = \{u = (u_1, \dots, u_d) \in I^d; (u_{(i-1)}, u_{(i)}) \subset \Psi_i^\delta \text{ for all } 2 \leq i \leq d\}.$$

We have the following Lemma for all absolutely continuous copula $C \in \mathcal{C}_\delta^0$ with density c .

Lemma 4.2. *Let $\delta \in \mathcal{D}^0$ and $2 \leq i \leq d$. Then for all copulas $C \in \mathcal{C}_\delta^0$ with density c , we have $c \mathbf{1}_{I \setminus L_\delta} = 0$ a.e., that is for a.e. $u = (u_1, \dots, u_d) \in I^d$, for all $s \notin \Psi_i^\delta$:*

$$(42) \quad c(u) \mathbf{1}_{\{u_{(i-1)} < s < u_{(i)}\}} = 0.$$

Proof. The complementary set $(\Psi_i^\delta)^c$ is given by:

$$(43) \quad (\Psi_i^\delta)^c = \bigcup_{j \in J_i} \overline{\{g_i^{(j)}, d_i^{(j)}\}}.$$

Let $U = (U_1, \dots, U_d)$ be a random vector with cdf $C \in \mathcal{C}_\delta^0$. For $2 \leq i \leq d$ and $s \in \bigcup_{j \in J_i} \{g_i^{(j)}, d_i^{(j)}\}$, that is $\delta_{(i-1)}(s) = \delta_{(i)}(s)$, we have:

$$\mathbb{P}(U_{(i-1)} < s < U_{(i)}) = \mathbb{P}(U_{(i-1)} < s) - \mathbb{P}(U_{(i)} \leq s) = \delta_{(i-1)}(s) - \delta_{(i)}(s) = 0.$$

This implies that (42) holds a.e. for all $s \in \bigcup_{j \in J_i} \{g_i^{(j)}, d_i^{(j)}\}$. Since J_i is at most countable, we have for a.e. $u \in I^d$ and for all $s \in \bigcup_{j \in J_i} \{g_i^{(j)}, d_i^{(j)}\}$, that (42) holds. Since for all $u \in I$, $s \notin \Psi_i^\delta$ there exists $s' \in \bigcup_{j \in J_i} \{g_i^{(j)}, d_i^{(j)}\}$ such that

$$\mathbf{1}_{\{u_{(i-1)} < s < u_{(i)}\}} = \mathbf{1}_{\{u_{(i-1)} < s' < u_{(i)}\}},$$

we can conclude that for a.e. $u \in I^d$ and for all $s \notin \Psi_i^\delta$ (42) hold. $\square$

Notice that for all $u = (u_1, \dots, u_d) \in I^d$:

$$(44) \quad \mathbf{1}_{L_\delta}(u) \leq \prod_{i=1}^d \mathbf{1}_{\Psi_i^\delta \cap \Psi_{i+1}^\delta}(u_{(i)}).$$

We define the function c_δ on I^d as, for $u = (u_1, \dots, u_d) \in I^d$:

$$(45) \quad c_\delta(u) = \frac{1}{d!} \mathbf{1}_{L_\delta}(u) \prod_{i=1}^d a_i(u_{(i)}),$$

where the function a_i , $1 \leq i \leq d$, are given by, for $t \in I$:

$$(46) \quad a_i(t) = K_i'(t) e^{K_{i+1}(t) - K_i(t)} \mathbf{1}_{\Psi_i^\delta \cap \Psi_{i+1}^\delta}(t),$$

with for $1 \leq i \leq d$, $t \in (g_i^{(j)}, d_i^{(j)})$:

$$(47) \quad K_i(t) = \int_{m_i^{(j)}}^t \frac{\delta'_{(i)}(s)}{\delta_{(i-1)}(s) - \delta_{(i)}(s)} ds$$

and the conventions $\delta_{(0)} = 1$ and $K_{d+1} = 0$. Notice that for $t \in \Psi_1^\delta$:

$$(48) \quad K_1(t) = -\log(1 - \delta_{(1)}(t)).$$

Remark 4.3. The choice of $m_i^{(j)}$ for the integration lower bound in (47) is arbitrary as any other value in $(g_i^{(j)}, d_i^{(j)})$ would not change the definition of c_δ in (45).

Remark 4.4. For all $1 \leq i \leq d$, $j \in J_i$, $t \in (m_i^{(j)}, d_i^{(j)})$, we have the following lower bound for $K_i(t)$:

$$K_i(t) \geq \int_{m_i^{(j)}}^t \frac{\delta'_{(i)}(s)}{\delta_{(i-1)}(d_i^{(j)}) - \delta_{(i)}(s)} ds = \log \left(\frac{\delta_{(i-1)}(d_i^{(j)}) - \delta_{(i)}(m_i^{(j)})}{\delta_{(i-1)}(d_i^{(j)}) - \delta_{(i)}(t)} \right).$$

Since $\delta_{(i)}$ is non-decreasing and $\delta_{(i-1)}(d_i^{(j)}) = \delta_{(i)}(d_i^{(j)})$, we have $\lim_{t \nearrow d_i^{(j)}} K_i(t) = +\infty$.

The following Proposition states that c_δ is the density of an absolutely continuous symmetric copula $C_\delta \in \mathcal{C}_\delta^0 \cap \mathcal{C}^{sym}$. It is more general than the results in [6], where only the diagonal $\delta_{(d)}$ was supposed given.

Proposition 4.5. *Let $\delta \in \mathcal{D}^0$. The function c_δ defined in (45)-(47) is the density of a symmetric copula $C_\delta \in \mathcal{C}_\delta^0 \cap \mathcal{C}^{sym}$. In addition, we have:*

$$(49) \quad H(C_\delta) = -\mathbb{J}(\delta) + \log(d!) + (d-1) + \sum_{i=1}^d H(\delta_{(i)}).$$

The proof of this Proposition is given in Section 6.2. The following characterization of C_δ is proved in Section 6.6.

Proposition 4.6. *Let $\delta \in \mathcal{D}^0$. Then C_δ is the only copula in $\mathcal{C}_\delta^0$ whose density is of the form $(1/d!) \mathbf{1}_{L_\delta}(u) \prod_{i=1}^d h_i(u_{(i)})$, where h_i , $1 \leq i \leq d$ are measurable non-negative functions defined on I .*

The following Theorem states that the unique optimal solution of $\max_{C \in \mathcal{C}_\delta} (H(C))$, if it exists, is given by C_δ . Its proof is given in Sections 6.7 for case (a) and 6.8 for case (b).

Theorem 4.7. *Let $\delta \in \mathcal{D}$.*

- (a) *If $\mathbb{J}(\delta) = +\infty$ then $\max_{C \in \mathcal{C}_\delta} H(C) = -\infty$.*
- (b) *If $\mathbb{J}(\delta) < +\infty$ then $\delta \in \mathcal{D}^0$, $\max_{C \in \mathcal{C}_\delta} H(C) > -\infty$ and C_δ given in Proposition 4.5 is the unique copula such that $H(C_\delta) = \max_{C \in \mathcal{C}_\delta} H(C)$.*

The copula C_δ will be called the maximum entropy copula with given multidagonal.

5. MAXIMUM ENTROPY DISTRIBUTION OF ORDER STATISTICS WITH GIVEN MARGINALS

We use the results of Section 4 to compute the density of the maximum entropy copula for marginals $\mathbf{F} \in \mathcal{F}_d^0$ with $\mathcal{F}_d^0$ defined in Section 3.3. Recall $\delta^\mathbf{F} = (\delta_{(1)}, \dots, \delta_{(d)}) = \mathbf{F} \circ G^{-1}$ and the definition of $\Sigma^{\delta^\mathbf{F}}$ in (31). Recall K_i defined by (47), for $1 \leq i \leq d$ and $T^\mathbf{F}$ defined by (34). We define the function $c_\mathbf{F}$ on I^d , for $u = (u_1, \dots, u_d) \in I^d$:

$$(50) \quad c_\mathbf{F}(u) = \prod_{i=2}^d \frac{e^{K_i(\delta_{(i-1)}^{-1}(u_{i-1})) - K_i(\delta_{(i)}^{-1}(u_i))}}{\delta_{(i-1)} \circ \delta_{(i)}^{-1}(u_i) - u_i} \mathbf{1}_{\{u \in T^\mathbf{F}; (\delta_{(1)}^{-1}(u_1), \dots, \delta_{(d)}^{-1}(u_d)) \in L_{\delta^\mathbf{F}}\}} \mathbf{1}_{\{\prod_{i=1}^d \delta'_{(i)} \circ \delta_{(i)}^{-1}(u_i) > 0\}}.$$

Recall the function $\mathbb{J}(\delta)$ defined on the set of multidagonals by (16) and $C_{\delta^\mathbf{F}}$ the copula with density given by (45)-(47).

Proposition 5.1. *Let $\mathbf{F} \in \mathcal{F}_d^0$. The function $c_\mathbf{F}$ defined by (50) is the density of the copula $C_\mathbf{F} = \mathcal{S}_\mathbf{F}^{-1}(C_{\delta^\mathbf{F}})$ which belongs to $\mathcal{C}^{OS}(\mathbf{F})$. The entropy of $C_\mathbf{F}$ is given by:*

$$(51) \quad H(C_\mathbf{F}) = d - 1 - \mathbb{J}(\delta^\mathbf{F}).$$

Proof. Since $\mathbf{F} \in \mathcal{F}_d^0$, we have that $\delta^\mathbf{F} \in \mathcal{D}^0$. According to Proposition 4.5, $c_{\delta^\mathbf{F}}$ defined by (45) is the density of a symmetric copula $C_{\delta^\mathbf{F}}$ which belongs to $\mathcal{C}^{sym}(\mathbf{F}) \cap \mathcal{C}^0$, thanks to Proposition 3.13 and Lemma 3.19. According to Lemma 3.19, formula (36) we get that $c_\mathbf{F} = s_\mathbf{F}^{-1}(C_{\delta^\mathbf{F}})$ is therefore the density of a copula $C_\mathbf{F}$ which belongs to $\mathcal{C}^{OS}(\mathbf{F}) \cap \mathcal{C}^0$. Use (36) and (7) to check (50). To conclude, use (38) and (49) to get (51). $\square$

Analogously to Lemma 4.2, we have the following restriction on the support of all $F \in \mathcal{L}_d^{OS}(\mathbf{F}) \cap \mathcal{L}_d^0$. Recall the definition of $\Psi_i^\mathbf{F}$ in (30). The proof of the next Lemma is similar to the proof of Lemma 4.2 and is left to the reader.

Lemma 5.2. *Let $\mathbf{F} \in \mathcal{F}_d^0$ and $t \in \mathbf{F}_i((\Psi_i^\mathbf{F})^c)$ for some $2 \leq i \leq d$. Then we have for all $F \in \mathcal{L}_d^{OS}(\mathbf{F}) \cap \mathcal{L}_d^0$ with density function f :*

$$f(x) \mathbf{1}_{\{x_{i-1} < t < x_i\}} = 0 \quad \text{for a.e. } x = (x_1, \dots, x_d) \in S.$$

For $\delta \in \mathcal{D}$, recall the definition of L_δ in (41). Let $L^\delta = L_\delta \cap S$. More generally, for $\mathbf{F} \in \mathcal{F}_d$, we set:

$$(52) \quad L^\mathbf{F} = \{x = (x_1, \dots, x_d) \in S; (x_{i-1}, x_i) \subset \Psi_i^\mathbf{F} \text{ for all } 2 \leq i \leq d\}.$$

The next Lemma establishes the connection between the sets $L_{\delta^\mathbf{F}}$ defined by (41) and $L^\mathbf{F}$.

Lemma 5.3. *Let $\mathbf{F} = (\mathbf{F}_1, \dots, \mathbf{F}_d) \in \mathcal{F}_d^0$ with density functions $\mathbf{f}_i$ for $1 \leq i \leq d$. Let $\delta^{\mathbf{F}} = (\delta_{(1)}, \dots, \delta_{(d)})$ given by (28), $T^{\mathbf{F}}$ given by (34) and $L_{\delta^{\mathbf{F}}}$ defined by (41). Then for $\prod_{i=1}^d \mathbf{f}_i(x_i) dx_1 \dots dx_d$ -a.e. $x \in \mathbb{R}^d$ we have that*

$$\mathbf{1}_{T^{\mathbf{F}}}(\mathbf{F}_1(x_1), \dots, \mathbf{F}_d(x_d)) \mathbf{1}_{L_{\delta^{\mathbf{F}}}}(\delta_{(1)}^{-1} \circ \mathbf{F}_1(x_1), \dots, \delta_{(d)}^{-1} \circ \mathbf{F}_d(x_d)) = \mathbf{1}_{L^{\mathbf{F}}}(x)$$

Proof. According to (4) and (6), we have $\mathbf{f}_i(t) dt$ -a.e. that $\mathbf{F}_i^{-1} \circ \mathbf{F}_i(t) = t$. This implies that $\prod_{i=1}^d \mathbf{f}_i(x_i) dx_1 \dots dx_d$ -a.e., $(\mathbf{F}_1(x_1), \dots, \mathbf{F}_d(x_d))$ belongs to $T^{\mathbf{F}}$ if and only if $x \in S$. Recall the sets $\Psi_i^{\delta^{\mathbf{F}}}$ given by (30). For $\prod_{i=1}^d \mathbf{f}_i(x_i) dx_1 \dots dx_d$ -a.e. $x \in S$, we have:

$$\begin{aligned} & (\delta_{(1)}^{-1} \circ \mathbf{F}_1(x_1), \dots, \delta_{(d)}^{-1} \circ \mathbf{F}_d(x_d)) \in L_{\delta^{\mathbf{F}}} \\ \iff & (\delta_{(i-1)}^{-1} \circ \mathbf{F}_{i-1}(x_{i-1}), \delta_{(i)}^{-1} \circ \mathbf{F}_i(x_i)) \subset \Psi_i^{\delta^{\mathbf{F}}}, \quad 2 \leq i \leq d \\ \iff & \forall t \in (\delta_{(i-1)}^{-1} \circ \mathbf{F}_{i-1}(x_{i-1}), \delta_{(i)}^{-1} \circ \mathbf{F}_i(x_i)) : \delta_{(i-1)}(t) > \delta_{(i)}(t), \quad 2 \leq i \leq d \\ \iff & \forall t \in (G \circ \mathbf{F}_{i-1}^{-1} \circ \mathbf{F}_{i-1}(x_{i-1}), G \circ \mathbf{F}_i^{-1} \circ \mathbf{F}_i(x_i)) : \mathbf{F}_{i-1} \circ G^{-1}(t) > \mathbf{F}_i \circ G^{-1}(t), \quad 2 \leq i \leq d \\ \iff & \forall t \in (G(x_{i-1}), G(x_i)) : \mathbf{F}_{i-1} \circ G^{-1}(t) > \mathbf{F}_i \circ G^{-1}(t), \quad 2 \leq i \leq d, \end{aligned}$$

where the first equivalence comes from the definition of $L_{\delta^{\mathbf{F}}}$, the second from the definition of $\Psi_i^{\delta^{\mathbf{F}}}$, the third from (28) and the last from the fact that $\mathbf{f}_i(t) dt$ -a.e. $\mathbf{F}_i^{-1} \circ \mathbf{F}_i(t) = t$. Consider the change of variable $s = G^{-1}(t)$. We have by (2):

$$t < G(x_i) \iff G^{-1}(t) < x_i \iff s < x_i.$$

Since $x_{i-1} \in I_g(\mathbf{F}_{i-1})$ $\mathbf{f}_{i-1}(x_{i-1}) dx_{i-1}$ -a.e., we get $x_{i-1} \in I_g(G)$ and by (6):

$$G(x_{i-1}) < t \iff x_{i-1} < G^{-1}(t) \iff x_{i-1} < s.$$

Therefore we deduce that $\prod_{i=1}^d \mathbf{f}_i(x_i) dx_1 \dots dx_d$ -a.e. $x \in S$:

$$\begin{aligned} & (\delta_{(1)}^{-1} \circ \mathbf{F}_1(x_1), \dots, \delta_{(d)}^{-1} \circ \mathbf{F}_d(x_d)) \in L_{\delta^{\mathbf{F}}} \\ \iff & \forall s \in (x_{i-1}, x_i) : \mathbf{F}_{i-1}(s) > \mathbf{F}_i(s), \quad 2 \leq i \leq d \\ \iff & x \in L^{\mathbf{F}}. \end{aligned}$$

□

Using Proposition 3.21, we check that the copula $C_{\mathbf{F}}$ maximizes the entropy over the set $\mathcal{C}^{OS}(\mathbf{F})$. We set for $x = (x_1, \dots, x_d) \in \mathbb{R}^d$:

$$(53) \quad F_{\mathbf{F}}(x) = C_{\mathbf{F}}(\mathbf{F}_1(x_1), \dots, \mathbf{F}_d(x_d)).$$

Let $\mathbf{f}_i$ denote the density function of $\mathbf{F}_i$ when it exists. Let us further note for $2 \leq i \leq d$, $t \in \mathbb{R}$:

$$(54) \quad \ell_i(t) = \frac{\mathbf{f}_i(t)}{\mathbf{F}_{i-1}(t) - \mathbf{F}_i(t)}.$$

When the densities $\mathbf{f}_i$ exist for all $1 \leq i \leq d$, we define the function $f_{\mathbf{F}}$ for $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ as:

$$(55) \quad f_{\mathbf{F}}(x) = \mathbf{f}_1(x_1) \prod_{i=2}^d \ell_i(x_i) \exp \left(- \int_{x_{i-1}}^{x_i} \ell_i(s) ds \right) \mathbf{1}_{L^{\mathbf{F}}}(x),$$

with $L^{\mathbf{F}}$ given by (52). The next theorem asserts that the cdf $F_{\mathbf{F}}$ maximizes the entropy over the set $\mathcal{L}_d^{OS}(\mathbf{F})$ and that its density is $f_{\mathbf{F}}$. Recall $\mathbb{J}$ defined by (16).

Theorem 5.4. *Let $\mathbf{F} = (\mathbf{F}_i, 1 \leq i \leq d) \in \mathcal{F}_d$.*

- (a) *If there exists $1 \leq i \leq d$ such that $H(\mathbf{F}_i) = -\infty$, or if $\mathbb{J}(\mathbf{F}) = +\infty$, then we have $\max_{F \in \mathcal{L}_d^{OS}(\mathbf{F})} H(F) = -\infty$.*
- (b) *If $H(\mathbf{F}_i) > -\infty$ for all $1 \leq i \leq d$, and $\mathbb{J}(\mathbf{F}) < +\infty$, then we have $\mathbf{F} \in \mathcal{F}_d^0$, $\max_{F \in \mathcal{L}_d^{OS}(\mathbf{F})} H(F) > -\infty$, and $F_{\mathbf{F}}$ defined in (53) is the unique cdf in $\mathcal{L}_d^{OS}(\mathbf{F})$ such that $H(F_{\mathbf{F}}) = \max_{F \in \mathcal{L}_d^{OS}(\mathbf{F})} H(F)$. Furthermore, the density function of $F_{\mathbf{F}}$ exists, and is given by $f_{\mathbf{F}}$ defined in (55). We also have:*

$$H(F_{\mathbf{F}}) = d - 1 + \sum_{i=1}^d H(\mathbf{F}_i) - \mathbb{J}(\mathbf{F}).$$

Proof. The proof of case (a) is postponed to Section 6.7.

We shall assume that $H(\mathbf{F}_i) > -\infty$ for all $1 \leq i \leq d$ and $\mathbb{J}(\delta^{\mathbf{F}}) < +\infty$. This implies that the densities $\mathbf{f}_i$ of $\mathbf{F}_i$ exist for $1 \leq i \leq d$ and, thanks to Lemma 3.17, that $\mathbf{F} \in \mathcal{F}_d^0$. Let $F_{\mathbf{F}}$ be defined by (53), that is the cdf with copula $C_{\mathbf{F}}$ from Proposition 5.1 and one-dimensional marginals cdf $\mathbf{F}$. Thanks to Proposition 5.1, we have $F_{\mathbf{F}} \in \mathcal{L}_d^{OS}(\mathbf{F})$.

We deduce from (15), Propositions 3.13 and 3.21, Theorem 4.7 case (b) and Proposition 5.1 that $F_{\mathbf{F}}$ is the only cdf such that $H(F_{\mathbf{F}}) = \max_{F \in \mathcal{L}_d^{OS}(\mathbf{F})} H(F)$. We deduce from (15), (51) and Lemma 3.14 that:

$$H(F_{\mathbf{F}}) = d - 1 + \sum_{i=1}^d H(\mathbf{F}_i) - \mathbb{J}(\mathbf{F}).$$

Since the copula $C_{\mathbf{F}}$ is absolutely continuous with density $c_{\mathbf{F}}$ given in (50), we deduce from (53) that $F_{\mathbf{F}}$ has density $f_{\mathbf{F}}$ given by, for a.e. $x = (x_1, \dots, x_d) \in \mathbb{R}^d$:

$$(56) \quad f_{\mathbf{F}}(x) = c_{\mathbf{F}}(\mathbf{F}_1(x_1), \dots, \mathbf{F}_d(x_d)) \prod_{i=1}^d \mathbf{f}_i(x_i).$$

Recall the expression (50) of $c_{\mathbf{F}}$ as well as K_i defined by (47), for $1 \leq i \leq d$. Using the change of variable $s = G^{-1}(t)$ and (24), we get (similarly to the proof of Lemma 3.14):

$$(57) \quad K_i \circ \delta_{(i)}^{-1} \circ \mathbf{F}_i(x_i) - K_i \circ \delta_{(i-1)}^{-1} \circ \mathbf{F}_{i-1}(x_{i-1}) = \int_{\mathbf{F}_{i-1}^{-1} \circ \mathbf{F}_{i-1}(x_{i-1})}^{\mathbf{F}_i^{-1} \circ \mathbf{F}_i(x_i)} \ell_i(s) ds.$$

Using (24), we also get:

$$(58) \quad \frac{\mathbf{f}_i(x_i)}{\delta_{(i-1)} \circ \delta_{(i)}^{-1} \circ \mathbf{F}_i(x_i) - \mathbf{F}_i(x_i)} = \frac{\mathbf{f}_i(x_i)}{\mathbf{F}_{i-1} \circ \mathbf{F}_i^{-1} \circ \mathbf{F}_i(x_i) - \mathbf{F}_i(x_i)}.$$

According to (6), we have $\mathbf{f}_i(t) dt$ -a.e. that $\mathbf{F}_i^{-1} \circ \mathbf{F}_i(t) = t$. For $1 \leq i \leq d$, we have from (6) that $\prod_{i=1}^d \mathbf{f}_i(x_i) dx_1 \cdots dx_d$ -a.e.:

$$(59) \quad \mathbf{1}_{\left\{ \prod_{i=1}^d \delta'_{(i)} \circ \delta_{(i)}^{-1}(\mathbf{F}_i(x_i)) > 0 \right\}} = 1.$$

We deduce from (50), (56), (57), (58), (59) and Lemma 5.3 that a.e. for $x = (x_1, \dots, x_d) \in \mathbb{R}^d$:

$$f_{\mathbf{F}}(x) = \mathbf{f}_1(x_1) \prod_{i=2}^d \ell_i(x_i) e^{-\int_{x_{i-1}}^{x_i} \ell_i(s) ds} \mathbf{1}_{L_{\mathbf{F}}}(x).$$

□

Remark 5.5. We deduce from the proof of Theorem 5.4 case (b) and Proposition 5.1, that if $\mathbf{F} = (\mathbf{F}_i, 1 \leq i \leq d) \in \mathcal{F}_d^0$, then $f_{\mathbf{F}}$ defined by (55) is a probability density function on $S \subset \mathbb{R}^d$.

Remark 5.6. The density $f_{\mathbf{F}}$ has a product form on $L^{\mathbf{F}}$, that is it can be written as, for a.e. $x = (x_1, \dots, x_d) \in \mathbb{R}^d$:

$$(60) \quad f_{\mathbf{F}}(x) = \prod_{i=1}^d p_i(x_i) \mathbf{1}_{L^{\mathbf{F}}}(x),$$

where the functions $(p_i, 1 \leq i \leq d)$ are measurable and non-negative.

In addition to Remark 5.6, the next Corollary asserts that $F_{\mathbf{F}}$ is the only element of $\mathcal{L}_d^{OS}(\mathbf{F})$, whose density has a product form.

Corollary 5.7. *Let $\mathbf{F} \in \mathcal{F}_d^0$. Let $F \in \mathcal{L}_d^{OS}(\mathbf{F})$ be an absolutely continuous cdf with density f given by, a.e. for $x = (x_1, \dots, x_d) \in \mathbb{R}^d$: $f(x) = \prod_{i=1}^d h_i(x_i) \mathbf{1}_{L^{\mathbf{F}}}(x)$, with $h_i, 1 \leq i \leq d$ some measurable non-negative functions on $\mathbb{R}$. Then we have $F = F_{\mathbf{F}}$ on $\mathbb{R}^d$.*

Proof. Let $X = (X_1, \dots, X_d)$ be an order statistic with cdf F , and F^{sym} the cdf of X_{Π} given by (17), with Π uniform on $\mathcal{S}_d$ and independent of X . Then the cdf F^{sym} is also absolutely continuous, and its density f^{sym} is given by :

$$(61) \quad f^{sym}(x) = \frac{1}{d!} \prod_{i=1}^d h_i(x_{(i)}) \mathbf{1}_{L^{\mathbf{F}}}(x^{OS}),$$

where x^{OS} is the ordered vector of x . The one-dimensional marginal cdf's of X_{Π} are all equal to G given by (18). Let $C \in \mathcal{C}^{OS}(\mathbf{F}) \cap \mathcal{C}^0$ denote the copula of F . Then according to (8), the copula $S_{\mathbf{F}}(C)$ of X_{Π} is given by, for a.e. $u = (u_1, \dots, u_d) \in I^d$:

$$S_{\mathbf{F}}(C)(u) = F^{sym}(G^{-1}(u)).$$

Therefore its density $s_{\mathbf{F}}(C)$ can be expressed as:

$$\begin{aligned} s_{\mathbf{F}}(C)(u) &= \frac{f^{sym}(G^{-1}(u))}{\prod_{i=1}^d g \circ G^{-1}(u_i)} \prod_{i=1}^d \mathbf{1}_{\{g \circ G^{-1}(u_i) > 0\}} \\ &= \frac{1}{d!} \mathbf{1}_{L^{\mathbf{F}}}(G^{-1}(u^{OS})) \prod_{i=1}^d \frac{h_i \circ G^{-1}(u_{(i)})}{g \circ G^{-1}(u_{(i)})} \mathbf{1}_{\{g \circ G^{-1}(u_{(i)}) > 0\}}, \end{aligned}$$

where g is the density of G . Notice that for $x = (x_1, \dots, x_d) \in S$, we have a.e.:

$$\mathbf{1}_{\{\prod_{i=1}^d h_i(x_i) > 0\}} \leq \mathbf{1}_{\{\prod_{i=1}^d f_i(x_i) > 0\}},$$

with f_i the density of X_i . Therefore by Lemma 5.3 and since G is continuous, we have that $\prod_{i=1}^d h_i \circ G^{-1}(u_{(i)}) du_1 \dots du_d$ -a.e.:

$$\mathbf{1}_{L^{\mathbf{F}}}(G^{-1}(u^{OS})) = \mathbf{1}_{L_{\delta \mathbf{F}}}(\delta_{(1)}^{-1} \circ \delta_{(1)}(u_{(1)}), \dots, \delta_{(d)}^{-1} \circ \delta_{(d)}(u_{(d)})).$$

By Lemma 3.12, $S_{\mathbf{F}}(C)$ belongs to $\mathcal{C}_{\delta \mathbf{F}}^0$ and thus $s_{\mathbf{F}}(C) = 0$ a.e. on $Z_{\delta \mathbf{F}}$ defined by (39). Then use (4) and (6) to get that $\delta_{(i)}^{-1} \circ \delta_{(i)}(u_{(i)}) = u_{(i)}$ a.e. on $Z_{\delta \mathbf{F}}^c$. This gives:

$$\mathbf{1}_{L^{\mathbf{F}}}(G^{-1}(u^{OS})) = \mathbf{1}_{L_{\delta \mathbf{F}}}(u^{OS}) = \mathbf{1}_{L_{\delta \mathbf{F}}}(u)$$

that is $s_{\mathbf{F}}(C)$ is of the form $s_{\mathbf{F}}(C)(u) = (1/d!) \mathbf{1}_{L_{\delta\mathbf{F}}}(u) \prod_{i=1}^d \bar{h}_i(u_{(i)})$ for some measurable non-negative functions $(\bar{h}_i, 1 \leq i \leq d)$. Then, thanks to Proposition 4.6, we get that $S_{\mathbf{F}}(C) = C_{\delta\mathbf{F}}$. Then, use Proposition 5.1 to get that $F = F_{\mathbf{F}}$. $\square$

6. PROOFS

6.1. Preliminary notations for the optimization problem. Recall notations from Sections 2 and 3. In particular if $u = (u_1, \dots, u_d) \in I^d$ then $u^{OS} = (u_{(1)}, \dots, u_{(d)})$ denote the ordered vector of u .

In order to apply the technique established in [5], we introduce the linear functional $\mathcal{A} = (\mathcal{A}_i, 1 \leq i \leq 2d) : L^1(I^d) \rightarrow L^1(I)^{2d}$ as, for $f \in L^1(I^d)$ and $r \in I$:

$$\mathcal{A}_i(f)(r) = \int_{I^d} f(u) \mathbf{1}_{\{u_i \leq r\}} du \quad \text{and} \quad \mathcal{A}_{d+i}(f)(r) = \int_{I^d} f(u) \mathbf{1}_{\{u_{(i)} \leq r\}} du \quad \text{for } 1 \leq i \leq d.$$

Let $\delta = (\delta_{(i)}, 1 \leq i \leq d) \in \mathcal{D}^0$ be a multidagonal, see Definition 3.6. We set $b^\delta = (b_i, 1 \leq i \leq 2d)$ given by $b_i = \text{id}_I$ the identity function on I and $b_{d+i} = \delta_{(i)}$, for $1 \leq i \leq d$. If, for $c \in L^1(I^d)$, we have $\mathcal{A}_i(c) = b_i, 1 \leq i \leq d$ and $c \geq 0$ a.e., then we deduce that c is the density of an absolutely continuous copula, say C . If we further have $\mathcal{A}_{d+i}(c) = b_{d+i}$, for $1 \leq i \leq d$, then δ is the multidagonal of C .

Lemma 6.1. *Let $\delta \in \mathcal{D}^0$ and $b^\delta = (b_i, 1 \leq i \leq 2d)$. If $c \in L^1(I^d)$ is non-negative, symmetric and satisfies $\mathcal{A}_{d+i}(c) = b_{d+i}$ for $1 \leq i \leq d$, then c is the density of a copula with multidagonal δ .*

Proof. The symmetry and non-negativity of c as well as the condition $\mathcal{A}_{d+1}(c)(1) = \int_{I^d} c = b_{d+1}(1) = 1$ ensures that c is a density function of an exchangeable random vector $V = (V_1, \dots, V_d)$ on I^d . Recall $V^{OS} = (V_{(1)}, \dots, V_{(d)})$ denotes the corresponding order statistics. By symmetry, the lemma is proved as soon as we check that $\mathcal{A}_1(c) = b_1$. We have for $r \in I$:

$$\mathcal{A}_1(c)(r) = \mathbb{P}(V_1 \leq r) = \sum_{i=1}^d \mathbb{P}(V_{(i)} \leq r | V_1 = V_{(i)}) \mathbb{P}(V_1 = V_{(i)}) = \sum_{i=1}^d \delta_{(i)}(r) \frac{1}{d} = r,$$

where we used the exchangeability of V and the definition of $\delta_{(i)}$ for the third equality, and (22) for the last. This gives $\mathcal{A}_1(c) = b_1$. $\square$

6.2. Proof of Proposition 4.5. Let $\delta \in \mathcal{D}^0$. Lemma 3.15 implies that $\delta_{(i)}((\Psi_i^\delta)^c)$ has zero Lebesgue measure for all $2 \leq i \leq d$ with Ψ_i^δ given by (30). By construction, the function c_δ defined by (45) is non-negative, symmetric and well defined a.e. on I^d . Recall the notation $(g_i^{(j)}, d_i^{(j)})$ used in (40). We define the functions B_i on I as, for $1 \leq i \leq d+1$, $t \in (g_i^{(j)}, d_i^{(j)})$ (with the conventions $\Psi_1^\delta = (0, d_1), \Psi_{d+1}^\delta = (g_{d+1}, 1)$):

$$(62) \quad B_{d+1}(t) = 1 \quad \text{and} \quad B_i(t) = \int_t^{d_i^{(j)}} a_i(s) B_{i+1}(s) ds \quad \text{for } 1 \leq i \leq d.$$

For $t \in (\Psi_i^\delta)^c$, we set $B_i(t) = 0$. Recall K_i defined in (47) for $1 \leq i \leq d+1$ with the convention $K_{d+1} = 0$. We show that B_i can be simply expressed by K_i on Ψ_i^δ .

Lemma 6.2. *Let $1 \leq i \leq d+1$ and $t \in \Psi_i^\delta$. Then we have:*

$$(63) \quad B_i(t) = \exp(-K_i(t)).$$

Proof. For $i = d + 1$, the result is trivial. We proceed by induction on i . We suppose that $B_{i+1}(t) = \exp(-K_{i+1}(t))$ holds for some $1 \leq i \leq d$, and all $t \in \psi_{i+1}^\delta$. We have for $t \in (g_i^{(j)}, d_i^{(j)})$:

$$\begin{aligned}
B_i(t) &= \int_t^{d_i^{(j)}} a_i(s) B_{i+1}(s) ds \\
&= \int_t^{d_i^{(j)}} K_i'(s) e^{K_{i+1}(s) - K_i(s)} \mathbf{1}_{\Psi_i^\delta \cap \Psi_{i+1}^\delta}(s) B_{i+1}(s) ds \\
&= \int_t^{d_i^{(j)}} K_i'(s) e^{-K_i(s)} \mathbf{1}_{\Psi_i^\delta \cap \Psi_{i+1}^\delta}(s) ds. \\
&= \int_t^{d_i^{(j)}} K_i'(s) e^{-K_i(s)} ds \\
&= \exp(-K_i(t)),
\end{aligned}$$

where we used the definition of a_i given by (46) for the second equality, the induction hypothesis for the third equality, $(t, d_i^{(j)}) \subset \Psi_i^\delta$ and Lemma 3.15 for the fourth equality, and finally Remark 4.4 for the fifth equality. This ends the induction. $\square$

Similarly, we define the functions E_i on I as, for $0 \leq i \leq d$ as for $t \in (g_{i+1}^{(j)}, d_{i+1}^{(j)})$:

$$(64) \quad E_0(t) = 1, \quad \text{and} \quad E_i(t) = \int_{g_{i+1}^{(j)}}^t a_i(s) E_{i-1}(s) ds \quad \text{for} \quad 1 \leq i \leq d.$$

For $t \in (\Psi_{i+1}^\delta)^c$ we set $E_i(t) = 0$. The next Lemma gives a simple formula for E_i on Ψ_{i+1}^δ .

Lemma 6.3. *Let $0 \leq i \leq d$ and $t \in \Psi_{i+1}^\delta$. Then we have:*

$$(65) \quad E_i(t) = (\delta_{(i)}(t) - \delta_{(i+1)}(t)) \exp(K_{i+1}(t)).$$

Proof. For $i = 0$ the result is clear thanks to the convention $\delta_{(0)} = 1$ and (48). We proceed by induction on i . We suppose that $E_{i-1}(t) = (\delta_{(i-1)}(t) - \delta_{(i)}(t)) \exp(K_i(t))$ holds for some $1 \leq i \leq d$, and all $t \in \psi_i^\delta$. Let us denote $h_i = \delta_{(i-1)} - \delta_{(i)}$. Before computing $E_i(t)$ for $t \in (g_{i+1}^{(j)}, d_{i+1}^{(j)})$, we give an alternative expression for $\exp(K_i(s))$ for $s \in (g_{i+1}^{(j)}, t)$:

$$\begin{aligned}
e^{K_{i+1}(s)} &= \exp \left(- \int_{m_{i+1}^{(j)}}^t \frac{h'_{i+1}(u)}{h_{i+1}(u)} + \int_s^t \frac{h'_{i+1}(u)}{h_{i+1}(u)} + \int_{m_{i+1}^{(j)}}^s \frac{\delta'_{(i)}(u)}{h_{i+1}(u)} du \right) \\
(66) \quad &= \frac{h_{i+1}(t)}{h_{i+1}(s)} \exp \left(- \int_{m_{i+1}^{(j)}}^t \frac{h'_{i+1}(u)}{h_{i+1}(u)} + \int_{m_{i+1}^{(j)}}^s \frac{\delta'_{(i)}(u)}{h_{i+1}(u)} du \right).
\end{aligned}$$

Then we have for $t \in (g_{i+1}^{(j)}, d_{i+1}^{(j)})$:

$$\begin{aligned}
 E_i(t) &= \int_{g_{i+1}^{(j)}}^t a_i(s) E_{i-1}(s) ds \\
 &= \int_{g_{i+1}^{(j)}}^t K'_i(s) e^{K_{i+1}(s)-K_i(s)} \mathbf{1}_{\Psi_i^\delta \cap \Psi_{i+1}^\delta}(s) E_{i-1}(s) ds \\
 &= \int_{g_{i+1}^{(j)}}^t K'_i(s) e^{K_{i+1}(s)} h_i(s) \mathbf{1}_{\Psi_i^\delta \cap \Psi_{i+1}^\delta}(s) ds \\
 &= \int_{g_{i+1}^{(j)}}^t \delta'_{(i)}(s) e^{K_{i+1}(s)} \mathbf{1}_{\Psi_i^\delta \cap \Psi_{i+1}^\delta}(s) ds \\
 &= h_{i+1}(t) \exp \left(- \int_{m_{i+1}^{(j)}}^t \frac{h'_{i+1}(u)}{h_{i+1}(u)} du \right) \int_{g_{i+1}^{(j)}}^t \left(\frac{\delta'_{(i)}(s)}{h_{i+1}(s)} \exp \left(\int_{m_{i+1}^{(j)}}^s \frac{\delta'_{(i)}(u)}{h_{i+1}(u)} du \right) \right) ds \\
 &= h_{i+1}(t) \exp \left(- \int_{m_{i+1}^{(j)}}^t \frac{h'_{i+1}(u) - \delta'_{(i)}(u)}{h_{i+1}(u)} du \right) \\
 &= h_{i+1}(t) \exp(K_{i+1}(t)),
 \end{aligned}$$

where we used the definition of a_i given by (46) for the second equality, the induction hypothesis for the third equality, Lemma 3.15 and (66) for the fifth equality, and for the seventh equality we use that, for $t \in (g_{i+1}^{(j)}, m_{i+1}^{(j)})$ (similarly to Remark 4.4):

$$\int_{m_{i+1}^{(j)}}^t \frac{\delta'_{(i)}(s)}{h_{i+1}(s)} ds \leq \int_{m_{i+1}^{(j)}}^t \frac{\delta'_{(i)}(s)}{\delta_{(i)}(s) - \delta_{(i+1)}(g_{i+1}^{(j)})} ds = \log \left(\frac{\delta_{(i)}(t) - \delta_{(i+1)}(g_{i+1}^{(j)})}{\delta_{(i)}(m_{i+1}^{(j)}) - \delta_{(i+1)}(g_{i+1}^{(j)})} \right),$$

giving $\lim_{t \searrow g_{i+1}^{(j)}} \int_{m_{i+1}^{(j)}}^t \frac{\delta'_{(i)}(s)}{h_{i+1}(s)} ds = -\infty$.

□

The following Lemma justifies the introduction of the functions B_i, E_i .

Lemma 6.4. *We have with $u_{(0)} = 0$ for $1 \leq i \leq d$, $t \in \Psi_i^\delta$:*

$$(67) \quad \int_{I^d} c_\delta(u) \mathbf{1}_{\{u_{(i-1)} \leq t \leq u_{(i)}\}} du = B_i(t) E_{i-1}(t).$$

Proof. The definition (62) of B_i for $1 \leq i \leq d$ gives that for $t \in I$:

$$B_i(t) = \int a_i(r_i) a_{i+1}(r_{i+1}) \dots a_d(r_d) \mathbf{1}_{\{t \leq r_i \leq r_{i+1} \leq \dots \leq r_d \leq 1\}} \mathbf{1}_{\{[t, r_i] \subset \Psi_i^\delta\}} \prod_{j=i}^{d-1} \mathbf{1}_{\{(r_j, r_{j+1}) \subset \Psi_{j+1}^\delta\}} dr,$$

with $r = (r_i, r_{i+1}, \dots, r_d) \in I^{d-i+1}$. Similarly, we have for $1 \leq i \leq d$, $t \in I$ that $E_{i-1}(t)$ is equal to:

$$\int a_1(q_1) a_2(q_2) \dots a_{i-1}(q_{i-1}) \mathbf{1}_{\{0 \leq q_1 \leq q_2 \leq \dots \leq q_{i-1} \leq t\}} \mathbf{1}_{\{[q_{i-1}, t] \subset \Psi_i^\delta\}} \prod_{j=1}^{i-2} \mathbf{1}_{\{(q_j, q_{j+1}) \subset \Psi_{j+1}^\delta\}} dq,$$

with $q = (q_1, q_2, \dots, q_{i-1}) \in I^{i-1}$. Multiplying $B_i(t)$ with $E_{i-1}(t)$ gives:

$$\begin{aligned} B_i(t)E_{i-1}(t) &= \int_{\Delta} \prod_{j=1}^d a_j(u_j) \mathbf{1}_{\{u_{i-1} \leq t \leq u_i\}} \prod_{j=1}^{d-1} \mathbf{1}_{\{(u_j, u_{j+1}) \subset \Psi_{j+1}^\delta\}} du \\ &= \int_{\Delta} \prod_{j=1}^d a_j(u_j) \mathbf{1}_{\{u_{i-1} \leq t \leq u_i\}} \mathbf{1}_{L^\delta}(u) du \\ &= d! \int_{\Delta} c_\delta(u) \mathbf{1}_{\{u_{i-1} \leq t \leq u_i\}} du \\ &= \int_{I^d} c_\delta(u) \mathbf{1}_{\{u_{(i-1)} \leq t \leq u_{(i)}\}} du, \end{aligned}$$

where we used the symmetry of c_δ for the fourth equality. $\square$

Lemma 6.4 with $i = 1$ ensures that $\int_{I^d} c_\delta(u) du = \lim_{t \searrow 0} B_1(t)E_0(t) = 1$, that is c_δ a probability density function on I^d . Now we compute $\mathcal{A}_{d+1}(c_\delta)$. We have, for $t \in \Psi_i^\delta$:

$$\mathcal{A}_{d+1}(c_\delta)(t) = \int_{I^d} c_\delta(u) \mathbf{1}_{\{u_{(1)} \leq t\}} du = 1 - \int_{I^d} c_\delta(u) \mathbf{1}_{\{u_{(1)} \geq t\}} du = 1 - B_1(t)E_0(t) = \delta_{(1)}(t),$$

where we used Lemma 6.4 with $i = 1$ for the third equality, then (63) and (48) for the fourth equality. By continuity this gives $\mathcal{A}_{d+1}(c_\delta) = \delta_{(1)}$ on I . For $2 \leq i \leq d$, we have by induction for $t \in \Psi_i^\delta$:

$$\begin{aligned} \mathcal{A}_{d+i}(c_\delta)(t) &= \int_{I^d} c_\delta(u) \mathbf{1}_{\{u_{(i)} \leq t\}} du \\ &= \int_{I^d} c_\delta(u) \mathbf{1}_{\{u_{(i-1)} \leq t\}} du - \int_{I^d} c_\delta(u) \mathbf{1}_{\{u_{(i-1)} \leq t \leq u_{(i)}\}} du \\ &= \mathcal{A}_{d+i-1}(c_\delta)(t) - B_i(t)E_{i-1}(t) \\ &= \delta_{(i-1)}(t) - (\delta_{(i-1)}(t) - \delta_{(i)}(t)) \\ &= \delta_{(i)}(t), \end{aligned}$$

where we used the induction and Lemma 6.4 for the third equality, as well as (63) and (65) for the fourth. By continuity, we obtain $\mathcal{A}_{d+i}(c_\delta) = \delta_{(i)}$ on I . Then use Lemma 6.1 to get that c_δ is the density of a (symmetric) copula, say C_δ , with multidagonal δ .

The computation of $H(C_\delta)$ is technical. By symmetry of C_δ , we have:

$$H(C_\delta) = -d! \int_{\Delta} c_\delta \log(c_\delta).$$

Let $\varepsilon \in (0, 1/2)$. We define for $1 \leq i \leq d$ the set $I_i^\varepsilon = \{t \in I; (t - \varepsilon, t + \varepsilon) \subset \Psi_i^\delta \cap \Psi_{i+1}^\delta\}$ as well as $I_0^\varepsilon = (\varepsilon, d_1 - \varepsilon)$ and $I_{d+1}^\varepsilon = (g_{d+1} + \varepsilon, 1 - \varepsilon)$. We set $\Delta^\varepsilon = \Delta \cap \prod_{i=1}^d I_i^\varepsilon$. Since $x \log(x) \geq -1/e$ for $x > 0$, we deduce by monotone convergence that:

$$(68) \quad H(C_\delta) = \lim_{\varepsilon \downarrow 0} H_\varepsilon(C_\delta),$$

with:

$$H_\varepsilon(C_\delta) = -d! \int_{\Delta^\varepsilon} c_\delta \log(c_\delta).$$

We can decompose $H_\varepsilon(C_\delta)$ as:

$$(69) \quad H_\varepsilon(C_\delta) = J_1(\varepsilon) + J_2(\varepsilon) + J_3(\varepsilon) + J_4(\varepsilon),$$

with:

$$\begin{aligned}
 J_1(\varepsilon) &= -d! \sum_{i=1}^d \int_{\Delta^\varepsilon} c_\delta(u) \log(\delta'_{(i)}(u_i)) du, \\
 J_2(\varepsilon) &= d! \sum_{i=2}^d \int_{\Delta^\varepsilon} c_\delta(u) \log(\delta_{(i-1)}(u_i) - \delta_{(i)}(u_i)) du, \\
 J_3(\varepsilon) &= d! \sum_{i=2}^d \int_{\Delta^\varepsilon} c_\delta(u) (K_i(u_i) - K_i(u_{i-1})) du, \\
 J_4(\varepsilon) &= \log(d!) d! \int_{\Delta^\varepsilon} c_\delta(u) du,
 \end{aligned}$$

where we used (48) to express $a_1 = \delta'_{(1)} e^{K_2}$ a.e., so that the sums in J_2 and J_3 start at $i = 2$.

Since $\mathcal{A}_{d+i}(c_\delta) = \delta_{(i)}$, we deduce that for $1 \leq i \leq d$ and any measurable non-negative function h defined on I :

$$(70) \quad \int_{I^d} c_\delta(u) h(u_{(i)}) du = \int_I \delta'_{(i)}(t) h(t) dt.$$

In particular, we get:

$$\int_{I^d} c_\delta(u) \left| \log(\delta'_{(i)}(u_{(i)})) \right| du = \int_I \delta'_{(i)}(t) \left| \log(\delta'_{(i)}(t)) \right| dt,$$

which is finite thanks to Remark 3.8. Therefore, by the dominated convergence and using (70) again, we have:

$$\lim_{\varepsilon \rightarrow 0} J_1(\varepsilon) = \sum_{i=1}^d H(\delta_{(i)}).$$

For $J_2(\varepsilon)$, we notice that the integrand is non-positive a.e., since for $t \in I$, $2 \leq i \leq d$, we have $\delta_{(i-1)}(t) - \delta_{(i)}(t) \leq 1$. Therefore, we get:

$$\lim_{\varepsilon \rightarrow 0} J_2(\varepsilon) = -d! \sum_{i=2}^d \int_{\Delta} c_\delta(u) \left| \log(\delta_{(i-1)}(u_i) - \delta_{(i)}(u_i)) \right| du = -\mathbb{J}(\delta),$$

where we used monotone convergence for the first equality and (70) as well as the definition (16) of $\mathbb{J}(\delta)$ for the second.

We now compute $J_3(\varepsilon)$. For $t \in I_{d+1}^\varepsilon$, we set $B_{d+1}^\varepsilon(t) = 1$, and for $t \in I_0^\varepsilon$, $E_0^\varepsilon(t) = 1$. For $1 \leq i \leq d$, we define for $t \in (g_i^{(j)}, d_i^{(j)})$ and $t' \in (g_{i+1}^{(j)}, d_{i+1}^{(j)})$:

$$B_i^\varepsilon(t) = \int_t^{\max(d_i^{(j)} - \varepsilon, t)} a_i(s) B_{i+1}^\varepsilon(s) ds, \quad \text{and} \quad E_i^\varepsilon(t') = \int_{\min(g_{i+1}^{(j)} + \varepsilon, t')}^{t'} a_i(s) E_{i-1}^\varepsilon(s) ds.$$

By monotone convergence, we have for $t \in \Psi_i^\delta$ and $t' \in \Psi_{i+1}^\delta$:

$$\lim_{\varepsilon \rightarrow 0} B_i^\varepsilon(t) = B_i(t) \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0} E_i^\varepsilon(t') = E_i(t').$$

An integration by parts gives:

$$\begin{aligned}
J_3(\varepsilon) &= d! \sum_{i=2}^d \left(\int_{\Delta^\varepsilon} c_\delta(u) K_i(u_i) du - \int_{\Delta^\varepsilon} c_\delta(u) K_i(u_{i-1}) du \right) \\
&= \sum_{i=2}^d \left(\int_{I_i^\varepsilon} E_{i-1}^\varepsilon(t) a_i(t) K_i(t) B_{i+1}^\varepsilon(t) dt - \int_{I_{i-1}^\varepsilon} E_{i-2}^\varepsilon(t) a_{i-1}(t) K_i(t) B_i^\varepsilon(t) dt \right) \\
&= \sum_{i=2}^d \left(\int_{I_i^\varepsilon} E_{i-1}^\varepsilon(t) a_i(t) K_i(t) B_{i+1}^\varepsilon(t) dt \right) \\
&\quad + \sum_{i=2}^d \left(\int_{I_i^\varepsilon} E_{i-1}^\varepsilon(t) K_i'(t) B_i^\varepsilon(t) dt - \int_{I_i^\varepsilon} E_{i-1}^\varepsilon(t) K_i(t) a_i(t) B_{i+1}^\varepsilon(t) dt \right) \\
&= \sum_{i=2}^d \int_{I_i^\varepsilon} E_{i-1}^\varepsilon(t) K_i'(t) B_i^\varepsilon(t) dt.
\end{aligned}$$

By monotone convergence and thanks to (47), (63) and (65), we get:

$$\lim_{\varepsilon \rightarrow 0} J_3(\varepsilon) = \sum_{i=2}^d \int_I E_{i-1}(t) K_i'(t) B_i(t) dt = \sum_{i=2}^d \int_I \delta'_{(i)}(t) dt = (d-1).$$

We deduce from (68) and (69) and the limits for J_1 , J_2 , J_3 and J_4 as ε goes down to 0 that:

$$H(C_\delta) = \sum_{i=1}^d H(\delta_{(i)}) - \mathbb{J}(\delta) + (d-1) + \log(d!).$$

6.3. The optimization problem. Let $\delta \in \mathcal{D}^0$. Recall notation from Section 6.1. The problem of maximizing H over $\mathcal{C}_\delta^0$ can be written as an optimization problem (P^δ) with infinite dimensional constraints:

$$(P^\delta) \quad \text{maximize } H(c) \text{ subject to } \begin{cases} \mathcal{A}(c) = b^\delta, \\ c \geq 0 \text{ a.e. and } c \in L^1(I^d). \end{cases}$$

Notice that if $f \in L^1(I^d)$ is non-negative and solves $\mathcal{A}(f) = b^\delta$, then f is the density of a copula. We say that a function f is feasible for (P^δ) if $f \in L^1(I^d)$, $f \geq 0$ a.e., $\mathcal{A}(f) = b^\delta$ and $H(f) > -\infty$. We say that f is an optimal solution of (P^δ) if f is feasible and $H(f) \geq H(g)$ for all g feasible. The next Proposition gives conditions which ensure the existence of an optimal solution.

Proposition 6.5. *Let $\delta \in \mathcal{D}^0$. If there exists c feasible for (P^δ) , then there exists a unique optimal solution to (P^δ) and it is symmetric.*

Proof. Since $\mathcal{A}(f) = b^\delta$ implies $\mathcal{A}_1(f)(1) = b_1(1)$ that is $\int_{I^d} f(x) dx = 1$, we can directly apply Corollary 2.3 of [5] which states that if there exists a feasible c , then there exists a unique optimal solution to (P^δ) . Since the constraints of (P^δ) are symmetric, such as the functional H , we deduce that if c^* is the optimal solution, then so is c_π^* defined for $\pi \in \mathcal{S}_d$ and $u \in I^d$ as $c_\pi^*(u) = c^*(u_\pi)$. By uniqueness of the optimal solution, we deduce that $c^* = c_\pi^*$ for all permutations $\pi \in \mathcal{S}_d$; hence c^* is symmetric. $\square$

Combining Lemmas 4.2 and 4.1 gives the following Corollary on the support of any c verifying $\mathcal{A}(c) = b^\delta$.

Corollary 6.6. *Let $\delta \in \mathcal{D}^0$. If $c \in L^1(I^d)$ is non-negative and verifies $\mathcal{A}(c) = b^\delta$, then $c = 0$ a.e. on $Z_\delta \cup L_\delta^c$ with L_δ defined by (41) and $L_\delta^c = I^d \setminus L_\delta$.*

6.4. Reduction of the optimization problem (P^δ) . Let $\delta \in \mathcal{D}^0$. Since the optimal solution of (P^δ) is symmetric, see Proposition 6.5, we can reduce the optimization problem by considering it on the simplex Δ . We define μ to be the Lebesgue measure restricted to $(Z_\delta^c \cap L_\delta) \cap \Delta$: $\mu(du) = \mathbf{1}_{(Z_\delta^c \cap L_\delta) \cap \Delta}(u)du$. We define, for $f \in L^1(I^d)$:

$$H^\mu(f) = - \int_{I^d} f(u) \log(f(u)) \mu(du).$$

From Corollary 6.6 we can deduce that if $c \in L^1(I^d)$ is non-negative symmetric and solves $\mathcal{A}(c) = b^\delta$, then:

$$(71) \quad H(c) = d! H^\mu(c).$$

Let us also define, for $f \in L^1(I^d)$, $1 \leq i \leq d$, $r \in I$:

$$\mathcal{A}_i^\mu(c)(r) = d! \int_{I^d} c(u) \mathbf{1}_{\{u_i \leq r\}} \mu(du).$$

We shall consider the restricted optimization problem (P_μ^δ) given by:

$$(P_\mu^\delta) \quad \text{maximize } H^\mu(c) \text{ subject to } \begin{cases} \mathcal{A}^\mu(c) = \delta, \\ c \geq 0 \text{ } \mu\text{-a.e. and } c \in L^1(I^d). \end{cases}$$

We have the following equivalence between (P^δ) and (P_μ^δ) . Recall u^{OS} denote the ordered vector of $u \in \mathbb{R}^d$.

Corollary 6.7. *Let $\delta \in \mathcal{D}^0$. If c is the optimal solution of (P^δ) then it is also an optimal solution to (P_μ^δ) . If $\hat{c}$ is an optimal solution of (P_μ^δ) , then c , defined by $c(u) = \hat{c}(u^{OS}) \mathbf{1}_{Z_\delta^c \cap L_\delta}(u)$ is the optimal solution to (P^δ) .*

Notice the Corollary implies that (P_μ^δ) has a μ -a.e. unique optimal solution: if c_1 and c_2 are two optimal solutions of (P_μ^δ) then μ -a.e. $c_1 = c_2$. Thanks to Proposition 6.5 and (71), Corollary 6.7 is a direct consequence of the following Lemma that establishes the connection between the constraints.

Lemma 6.8. *Let $\delta \in \mathcal{D}^0$. For $c \in L^1(I^d)$ symmetric and non-negative the following two conditions are equivalent:*

- (1) $\mathcal{A}(c) = b^\delta$.
- (2) $\mathcal{A}^\mu(c) = \delta$ and $c = 0$ a.e. on $Z_\delta \cup L_\delta^c$.

Proof. Assume that $\mathcal{A}(c) = b^\delta$. We have, by Corollary 6.6, that $c = 0$ a.e. on $Z_\delta \cup L_\delta^c$. This and the symmetry of c gives, for $1 \leq i \leq d$, $r \in I$:

$$\mathcal{A}_i^\mu(c)(r) = d! \int_{I^d} c(u) \mathbf{1}_{\{u_{(i)} \leq r\}} \mathbf{1}_\Delta(u) du = \int_{I^d} c(u) \mathbf{1}_{\{u_{(i)} \leq r\}} du = \delta_{(i)}(r).$$

On the other hand, let us assume that $\mathcal{A}^\mu(c) = \delta$ and $c = 0$ a.e. on $Z_\delta \cup L_\delta^c$. We have, for $1 \leq i \leq d$, $r \in I$:

$$\mathcal{A}_{d+i}(c)(r) = \int_{I^d} c(u) \mathbf{1}_{\{u_{(i)} \leq r\}} \mathbf{1}_{Z_\delta^c \cap L_\delta}(u) du = d! \int_{I^d} c(u) \mathbf{1}_{\{u_i \leq r\}} \mu(du) = \delta_{(i)}(r),$$

where we used $c = 0$ a.e. on $Z_\delta \cup L_\delta^c$ for the first equality, the symmetry of c and the definition of μ for the second, and $\mathcal{A}^\mu(c) = \delta$ for the third. Lemma 6.1 ensures then that $\mathcal{A}_i(c) = b_i$ for $1 \leq i \leq d$. This ends the proof. $\square$

6.5. Solution for the reduced optimization problem (P_μ^δ) . Let $\delta \in \mathcal{D}^0$. We compute $(\mathcal{A}^\mu)^* : L^\infty(I)^d \rightarrow L^\infty(I^d)$ the adjoint of $\mathcal{A}^\mu$. For $\lambda = (\lambda_i, 1 \leq i \leq d) \in L^\infty(I)^d$ and $f \in L^1(I^d)$, we have:

$$\langle (\mathcal{A}^\mu)^*(\lambda), f \rangle = \langle \lambda, \mathcal{A}^\mu(f) \rangle = \sum_{i=1}^d \int_I \lambda_i(r) \int_{I^d} f(u) \mathbf{1}_{\{u_i \leq r\}} d\mu(u) dr = \int_{I^d} f(u) \sum_{i=1}^d \Lambda_i(u_i) d\mu(u),$$

where we used the definition of the adjoint operator for the first equality, Fubini's theorem for the second, and the following definition of the functions $(\Lambda_i, 1 \leq i \leq d)$ for the third:

$$\Lambda_i(t) = \int_I \lambda_i(r) \mathbf{1}_{\{r \geq t\}} dr, \quad t \in I.$$

Thus, we have for $\lambda \in L^\infty(I)^d$ and $u = (u_1, \dots, u_d) \in I^d$:

$$(72) \quad (\mathcal{A}^\mu)^*(\lambda)(u) = \sum_{i=1}^d \Lambda_i(u_i).$$

We will use Theorem 2.9. from [5] on abstract entropy minimization, which we recall here, adapted to the context of (P_μ^δ) .

Theorem 6.9 (Borwein, Lewis and Nussbaum). *Suppose there exists $c > 0$ μ -a.e. which is feasible for (P_μ^δ) . Then there exists a μ -a.e. unique optimal solution, c^* , of (P_μ^δ) . Furthermore, we have $c^* > 0$ μ -a.e. and there exists a sequence $(\lambda^n, n \in \mathbb{N}^*)$ of elements of $L^\infty(I)^d$ such that:*

$$(73) \quad \int_{I^d} c^*(u) |(\mathcal{A}^\mu)^*(\lambda^n)(u) - \log(c^*(u))| \mu(du) \xrightarrow{n \rightarrow \infty} 0.$$

Now we are ready to prove that the optimal solution c^* of (P_μ^δ) is the product of measurable univariate functions.

Lemma 6.10. *Let $\delta \in \mathcal{D}^0$. Suppose that there exists $c > 0$ μ -a.e. which is feasible for (P_μ^δ) . Then there exist non-negative, measurable functions $(a_i^*, 1 \leq i \leq d)$ defined on I such that $a_i^*(s) = 0$ if $\delta'_{(i)}(s) = 0$ and the function c^* defined a.e. on I^d by:*

$$c^*(u) = \frac{1}{d!} \mathbf{1}_{L_\delta}(u) \prod_{i=1}^d a_i^*(u_i)$$

is the optimal solution to (P_μ^δ) .

Proof. According to Theorem 6.9, there exists a sequence $(\lambda^n, n \in \mathbb{N})$ of elements of $L^\infty(I)^d$ such that the optimal solution, say c^* , satisfies (73). This implies, thanks to (72), that there exist d sequences $(\Lambda_i^n, n \in \mathbb{N}^*, 1 \leq i \leq d)$ of elements of $L^\infty(I)$ such that the following convergence holds in $L^1(I^d, c^*\mu)$:

$$(74) \quad \sum_{i=1}^d \Lambda_i^n(u_i) \xrightarrow{n \rightarrow \infty} \log(c^*(u)).$$

We first assume that there exist Λ_i , $1 \leq i \leq d$ measurable functions defined on I such that μ -a.e. on S :

$$(75) \quad \sum_{i=1}^d \Lambda_i(u_i) = \log(c^*(u)).$$

Set $a_i^* = \sqrt[d]{d!} \exp(\Lambda_i)$ so that μ -a.e. on S :

$$(76) \quad c^*(u) = \frac{1}{d!} \prod_{i=1}^d a_i^*(u_i).$$

Recall $\mu(du) = \mathbf{1}_{(Z_\delta^c \cap L_\delta) \cap \Delta}(u) du$. From the definition (39) of Z_δ , we deduce that without loss of generality, we can assume that $a_i^*(u_i) = 0$ if $\delta'_{(i)}(u_i) = 0$. Therefore we obtain $c^*(u) = (1/d!) \mathbf{1}_{L_\delta}(u) \prod_{i=1}^d a_i^*(u_i)$ for $u \in I^d$.

To complete the proof, we now show that (75) holds for Λ_i , $1 \leq i \leq d$ measurable functions. We introduce the notation $u_{(-j)} = (u_1, \dots, u_{j-1}, u_{j+1}, \dots, u_d) \in I^{d-1}$. Let us define the probability measure $P(du) = c^*(u) \mu(du) / \int_{I^d} c^*(y) \mu(dy)$ on I^d . We fix j , $1 \leq j \leq d$. In order to apply Proposition 2 of [17], which ensures the existence of the limiting measurable functions Λ_i , $1 \leq i \leq d$, we first check that P is absolutely continuous with respect to $P_1^j \otimes P_2^j$, where $P_1^j(du_{(-j)}) = \int_{u_j \in I} P(du_{(-j)} du_j)$ and $P_2^j(du_j) = \int_{u_{(-j)} \in I^{d-1}} P(du_{(-j)} du_j)$ are the marginals of P . Notice that there exists a non-negative density function h such that $P(du) = h(u_{(-j)}, u_j) du_{(-j)} du_j$. Let $h_1(u_{(-j)}) = \int h(u_{(-j)}, u_j) du_j$ and $h_2(u_j) = \int h(u_{(-j)}, u_j) du_{(-j)}$ denote the density of the marginals P_1^j and P_2^j . Then the density of the product measure $P_1^j \otimes P_2^j$ is given by $P_1^j \otimes P_2^j(du) = h_1(u_{(-j)}) h_2(u_j) du_{(-j)} du_j$. The support of the density h is noted by $T_0 = \{u \in I^d; h(u) > 0\}$, and the support of the marginals are noted by $T_1 = \{v \in I^{d-1}; h_1(v) > 0\}$ and $T_2 = \{t \in I; h_2(t) > 0\}$. With this notation, we have that a.e. $T_0 \subset T_1 \times T_2$ (that is $T_0 \cap (T_1 \times T_2)^c$ is of zero Lebesgue measure). If $A \subset I^d$ is such that $\int \mathbf{1}_A(u) h_1(u_{(-j)}) h_2(u_j) du_{(-j)} du_j = 0$, then we also have $\int \mathbf{1}_{A \cap (T_1 \times T_2)}(u) h_1(u_{(-j)}) h_2(u_j) du_{(-j)} du_j = 0$. Since $h_1 h_2$ is positive on $T_1 \times T_2$, this implies that $A \cap (T_1 \times T_2)$ has zero Lebesgue measure. Therefore we have:

$$\int \mathbf{1}_A(u) h(u) du = \int \mathbf{1}_{A \cap (T_1 \times T_2)}(u) h(u) du + \int \mathbf{1}_{A \setminus (T_1 \times T_2)}(u) h(u) du = 0,$$

since $h = 0$ a.e. on $A \setminus (T_1 \times T_2)$. This proves that P is absolutely continuous with respect to $P_1^j \otimes P_2^j$. Then according to Proposition 2 of [17], (74) implies that there exist measurable functions Φ_j and $\tilde{\Lambda}_j$ defined respectively on I^{d-1} and I , such that c^* - μ -a.e. on Δ :

$$\log(c^*(u)) = \Phi_j(u_{(-j)}) + \tilde{\Lambda}_j(u_j).$$

As μ -a.e. $c^* > 0$, this equality holds μ -a.e. on S . Since we have such a representation for every $1 \leq j \leq d$, we can easily verify that $\log(c^*(u)) = \sum_{i=1}^d \Lambda_i(u_i)$ μ -a.e. with $\tilde{\Lambda}_j = \Lambda_j$ up to an additive constant. $\square$

6.6. Proof of Proposition 4.6. Let $\delta \in \mathcal{D}^0$. Recall that u^{OS} denotes the ordered vector of $u \in \mathbb{R}^d$. Let c be the density of a symmetric copula in $\mathbb{R}^d$ such that $\mathcal{A}(c) = b^\delta$ and c is of

product form, that is, thanks to Corollary 6.7, $c(u) = c^*(u^{OS})$ with

$$c^*(u) = \frac{1}{d!} \mathbf{1}_{L_\delta}(u) \prod_{i=1}^d a_i^*(u_i), \quad u = (u_1, \dots, u_d) \in \Delta,$$

where a_i^* , $1 \leq i \leq d$ are measurable non-negative functions defined on I . In this section, we shall prove that c equals c_δ defined by (45); that is, for all $1 \leq i \leq d$, a_i^* is a.e. equal, up to a multiplicative constant, to a_i defined in (46). This will prove Proposition 4.6.

Recall the definitions of $g_i^{(j)}$, $m_i^{(j)}$, $d_i^{(j)}$ from Section 4, for $1 \leq i \leq d+1$. We deduce from (44) that:

$$c^*(u) = \frac{1}{d!} \mathbf{1}_{L_\delta}(u) \prod_{i=1}^d a_i^*(u_i) \mathbf{1}_{\Psi_i^\delta \cap \Psi_{i+1}^\delta}(u_i), \quad u = (u_1, \dots, u_d) \in \Delta.$$

We deduce also from Lemma 6.8 that $\mathcal{A}^\mu(c^*) = \delta$. We introduce the following family of functions:

$$B_{d+1}^*(t) = E_0^*(t) = 1,$$

and for $1 \leq i \leq d$, $t \in (g_i^{(j)}, d_i^{(j)})$ and $t' \in (g_{i+1}^{(j)}, d_{i+1}^{(j)})$:

$$B_i^*(t) = \int_t^{d_i^{(j)}} a_i^*(s) B_{i+1}^*(s) ds, \quad E_i^*(t') = \int_{g_{i+1}^{(j)}}^{t'} a_i^*(s) E_{i-1}^*(s) ds.$$

Recall the functions B_i , for $1 \leq i \leq d+1$, and E_i , for $0 \leq i \leq d$ defined by (62) and (64). We will prove by (downward) induction on $i \in \{1, \dots, d+1\}$ that:

$$(77) \quad B_i^*(t) = B_i^*(m_i^{(j)}) B_i(t), \quad t \in (g_i^{(j)}, d_i^{(j)}).$$

For $i = d+1$, it trivially holds. Let us assume that (77) holds for $i+1$, $d \geq i \geq 1$. Recall the convention $K_{d+1} = 0$, $\delta_{(d+1)} = 0$ and $\delta_{(0)} = 1$. Arguing as in the proof of Lemma 6.4, we deduce from $\mathcal{A}_i^\mu(c^*) = \delta_{(i)}$ that for $r \in \Psi_i^\delta$:

$$\begin{aligned} \delta_{(i)}(r) &= d! \int_{I^d} c^*(u) \mathbf{1}_{\{u_{(i)} \leq r\}} \mu(du) \\ &= d! \int_{I^d} c^*(u) \mathbf{1}_{\{u_{(i+1)} \leq r\}} \mu(du) + \int_{\Delta} \mathbf{1}_{L_\delta}(u) \prod_{j=1}^d \left(a_j^*(u_j) \mathbf{1}_{\Psi_j^\delta \cap \Psi_{j+1}^\delta}(u_j) \right) \mathbf{1}_{\{u_i \leq r \leq u_{i+1}\}} du \\ &= \delta_{(i+1)}(r) + B_{i+1}^*(r) E_i^*(r). \end{aligned}$$

This gives on Ψ_i^δ :

$$(78) \quad \delta_{(i)} - \delta_{(i+1)} = B_{i+1}^* E_i^*.$$

Notice (78) holds for $i = d$ thanks to the conventions. We get on Ψ_i^δ :

$$\delta'_{(i)} - \delta'_{(i+1)} = -K'_{i+1} B_{i+1}^* E_i^* + B_{i+1}^* a_i^* E_{i-1}^* = -\delta'_{(i+1)} + B_{i+1}^* a_i^* E_{i-1}^*.$$

where we took the derivative in (78), twice the induction hypothesis for B_{i+1}^* and (63) for the first equality; then (47) and (78) for the second. We deduce that on Ψ_i^δ :

$$(79) \quad \delta'_{(i)} = B_{i+1}^* a_i^* E_{i-1}^*.$$

On Ψ_i^δ , we can divide (79) by (78) and get, thanks to (47):

$$\frac{a_i^* B_{i+1}^*}{B_i^*} = \frac{\delta'_{(i)}}{\delta_{(i-1)} - \delta_{(i)}} = K'_i.$$

Notice that $(B_i^*)' = -a_i^* B_{i+1}^*$. So using the representation (63) of B_i , we get that (77) holds for i . Thus (77) holds for $1 \leq i \leq d+1$. Then use (77) as well as $(B_i^*)' = -a_i^* B_{i+1}^*$ and $B'_i = -a_i B_{i+1}$ to get that for $t \in (g_i^{(j)}, d_i^{(j)}) \cap (g_{i+1}^{(k)}, d_{i+1}^{(k)})$:

$$a_i^*(t) = \frac{B_i^*(m_i^{(j)})}{B_{i+1}^*(m_{i+1}^{(k)})} a_i(t).$$

Therefore if $u = (u_1, \dots, u_d) \in L^\delta$, we have:

$$\prod_{i=1}^d a_i^*(u_{(i)}) = \frac{B_1^*(m_1)}{B_{d+1}^*(m_{d+1})} \prod_{i=1}^d a_i(u_{(i)}),$$

since when $u \in L_\delta$, $u_{(i-1)}$ and $u_{(i)}$ belong to the same interval $(g_i^{(j)}, d_i^{(j)})$ for $2 \leq i \leq d$. This ensures that c_δ and c^* are densities of probability function which differ by a multiplicative constant, therefore they are equal. This ends the proof of Proposition 4.6.

6.7. Proof of case (a) for Theorems 4.7 and 5.4. We first consider the case $d = 2$. Let $\delta \in \mathcal{D}^0$ with $\mathbb{J}(\delta) = +\infty$. Recall $\mathbb{J}(\delta)$ is defined by (16). We have:

$$\begin{aligned} \mathbb{J}(\delta) &= - \int_I \delta'_{(2)}(t) \log(2(t - \delta_{(2)}(t))) dt \\ &= -\log(2) - \int_I \log(t - \delta_{(2)}(t)) dt + \int_I (1 - \delta'_{(2)}(t)) \log(t - \delta_{(2)}(t)) dt \\ &= -\log(2) - \int_I \log(t - \delta_{(2)}(t)) dt + [(t - \delta_{(2)}(t)) \log(t - \delta_{(2)}(t)) - (t - \delta_{(2)}(t))]_0^1 \\ &= -\log(2) - \int_I \log(t - \delta_{(2)}(t)) dt, \end{aligned}$$

where we used $\delta_{(1)} + \delta_{(2)} = 2t$ for the first equality, $\delta_{(2)}(1) = 1$ and $\delta_{(2)}(0) = 0$ for the second and last. In particular, we obtain that $\mathbb{J}(\delta)$ is equal to $-\log(2) + \mathcal{J}(\delta_{(2)})$, with $\mathcal{J}$ as also defined by (1) in [6]. Therefore we deduce case (a) of Theorem 4.7 (for $d = 2$) from case (a) of Theorem 2.4 in [6]. Then, we get from (15) and Theorem 4.7 case (a) that $H(F) = -\infty$ for all $F \in \mathcal{L}_2^{OS}(\mathbf{F})$. This proves case (a) for Theorem 5.4 (for $d = 2$).

We then consider the case $d \geq 2$. Let $\delta \in \mathcal{D}^0$ with $\mathbb{J}(\delta) = +\infty$. This implies that there exists $2 \leq i \leq d$ such that $\int_I \delta'_{(i)}(t) |\log(\delta_{(i-1)}(t) - \delta_{(i)}(t))| dt = +\infty$. Set $\mathbf{F} = (\delta_{(i-1)}, \delta_{(i)})$ and notice that $\mathbf{F}$ belongs to $\mathcal{F}_2$ as $\delta_{(i)}$ is d -Lipschitz. Since $\int_I \delta'_{(i)}(t) |\log(\delta_{(i-1)}(t) - \delta_{(i)}(t))| dt = +\infty$, we deduce from the first part of this Section that $\max_{F \in \mathcal{L}_2^{OS}(\mathbf{F})} H(F) = -\infty$.

Consider a copula C belonging to $\mathcal{C}_\delta \cap \mathcal{C}^{sym}$ and U a random vector on I^d with cdf C . According to Lemma 3.20 and Lemma 3.3, as C is symmetric, we have:

$$H(U^{OS}) = H(U) - \log(d!) = H(C) - \log(d!).$$

It is easy to check that if $X = (X_1, \dots, X_d)$ is a random vector on I^d and $2 \leq i \leq d$, then we have $H((X_{i-1}, X_i)) \geq H(X)$. This implies that, for $V = (U_{i-1}^{OS}, U_i^{OS})$,

$$H(V) \geq H(C) - \log(d!).$$

Since the cdf of U_ℓ^{OS} is $\delta_{(\ell)}$ as $C \in \mathcal{C}_\delta$, we deduce the cdf of V belongs to $\mathcal{L}_2^{OS}(\mathbf{F})$, and thus $H(V) = -\infty$. This implies that $H(C) = -\infty$. Thanks to Proposition 6.5 which states that the entropy is maximal on symmetric copulas, we deduce that:

$$\max_{C \in \mathcal{C}_\delta} H(C) = \max_{C \in \mathcal{C}_\delta \cap \mathcal{C}^{sym}} H(C) = -\infty.$$

This proves cases (a) for Theorem 4.7. Then, we get from (15) that $H(F) = -\infty$ for all $F \in \mathcal{L}_d^{OS}(\mathbf{F})$. This proves case (a) for Theorem 5.4.

6.8. Proof of Theorem 4.7, case (b). Let $\delta \in \mathcal{D}$ with $\mathbb{J}(\delta) < +\infty$. Thanks to Lemma 3.15, $\mathbb{J}(\delta) < +\infty$ implies that $\delta \in \mathcal{D}^0$. By construction, c_δ introduced in Proposition 4.5 verifies $\mu - a.e.$ $c_\delta > 0$. The density c_δ is a feasible solution to the problem (P_μ^δ) . Theorem 6.9 ensures the existence of a unique optimal solution c^* . Furthermore, by Lemma 6.10, we have that there exist non-negative, measurable functions a_i^* , $1 \leq i \leq d$, such that $c^*(u) = (1/d!) \mathbf{1}_{L_\delta}(u) \prod_{i=1}^d a_i^*(u_i)$ μ -a.e. By Corollary 6.7, the optimal solution c of (P^δ) is given by, for $u = (u_1, \dots, u_d)$:

$$c(u) = c^*(u^{OS}) \mathbf{1}_{Z_\delta^c \cap L_\delta}(u) = \frac{1}{d!} \mathbf{1}_{L_\delta}(u) \prod_{i=1}^d a_i^*(u_{(i)}) \mathbf{1}_{\{\delta'_{(i)}(u_{(i)}) \neq 0\}}.$$

Since c is of product form, Proposition 4.6 yields that $c = c_\delta$ a.e., therefore C_δ is the unique copula achieving $H(C_\delta) = \max_{C \in \mathcal{C}_\delta} H(C)$.

7. OVERVIEW OF THE NOTATIONS

- $\mathcal{F}_d$: set of continuous one-dimensional marginals cdf $\mathbf{F} = (\mathbf{F}_1, \dots, \mathbf{F}_d)$ of d -dimensional order statistics, see (11).
- $\mathcal{F}_d^0$: set of continuous one-dimensional marginals cdf $\mathbf{F} = (\mathbf{F}_1, \dots, \mathbf{F}_d)$ of d -dimensional abs. cont. order statistics, see Definition 3.16.
- $\mathcal{L}_d$: set of all cdf's on $\mathbb{R}^d$.
- $\mathcal{L}_d^{1c}$: set of cdf's on $\mathbb{R}^d$ with continuous one-dimensional marginals cdf.
- $\mathcal{L}_d^0$: set of absolutely continuous cdf's on $\mathbb{R}^d$.
- $\mathcal{L}_d^{OS}$: set of cdf's of d -dimensional order statistics with continuous one dimensional marginals cdf.
- $\mathcal{L}_d^{OS}(\mathbf{F})$: set of cdf's of d -dimensional order statistics with marginals cdf $\mathbf{F}$, see (12).
- $\mathcal{L}_d^{sym}$: set of symmetric cdf's on $\mathbb{R}^d$.
- $S_{\mathbf{F}}$: symmetrizing operator on copulas, associated to the marginals cdf $\mathbf{F}$, see Definition 3.1.
- $\mathcal{C}$: set of all copulas.
- $\mathcal{C}^0$: set of absolutely continuous copulas.
- $\mathcal{C}^{OS}(\mathbf{F})$: set of copulas of order statistics with marginals cdf $\mathbf{F}$, see (13).
- $\mathcal{C}^{sym}$: set of symmetric (permutation invariant) copulas.
- $\mathcal{C}^{sym}(\mathbf{F})$: image of the set $\mathcal{C}^{OS}(\mathbf{F})$ by the operator $S_{\mathbf{F}}$, see (20). It is the set of symmetric copulas with multidagonal $\delta^{\mathbf{F}}$.
- $\mathcal{C}_{\delta}$: set of copulas with multidagonal δ , see Section 3.2.
- $\mathcal{C}_{\delta}^0$: set of abs. cont. copulas with multidagonal δ , see Section 3.2.
- $\mathcal{D}$: set of multidagonals of copulas, see Section 3.2.
- $\mathcal{D}^0$: set of multidagonals of abs. cont. copulas, see Section 3.2.
- $\Psi_i^{\mathbf{F}}$: set of points $t \in \mathbb{R}$ for which the marginals cdf $\mathbf{F} = (\mathbf{F}_1, \dots, \mathbf{F}_d)$ verify $\mathbf{F}_{i-1}(t) > \mathbf{F}_i(t)$, see (30).
- $T^{\mathbf{F}}$: set of points $u = (u_1, \dots, u_d) \in I^d$ for which $\mathbf{F}_1^{-1}(u_1) \leq \dots \leq \mathbf{F}_d^{-1}(u_d)$, see (34). The density of all copulas in $\mathcal{C}^{OS}(\mathbf{F})$ vanishes on $T^{\mathbf{F}}$.
- $L^{\mathbf{F}}$: set of ordered vectors $x \in \mathbb{R}^d$ such that the marginals cdf's $\mathbf{F} = (\mathbf{F}_1, \dots, \mathbf{F}_d)$ verify $\mathbf{F}_{i-1}(t) > \mathbf{F}_i(t)$ for all $t \in (x_{i-1}, x_i)$, $2 \leq i \leq d$, see (52). The density of any abs. cont. cdf in $\mathcal{L}_d^{OS}(\mathbf{F})$ vanishes outside $L^{\mathbf{F}}$.
- L_{δ} : set of points $u = (u_1, \dots, u_d) \in I^d$ for which all points $t \in (u_{(i-1)}, u_{(i)})$ verify $\delta_{(i-1)}(t) > \delta_{(i)}(t)$ for all $2 \leq i \leq d$, see (41). The density of any copula in $\mathcal{C}_{\delta}^0$ vanishes outside L_{δ} .
- Z_{δ} : set of points $u = (u_1, \dots, u_d) \in I^d$ such that $\delta'_{(i)}(u_{(i)}) = 0$ for some $1 \leq i \leq d$, see (39). The density of any copula in $\mathcal{C}_{\delta}^0$ vanishes on Z_{δ} .

REFERENCES

- [1] B. C. Arnold, N. Balakrishnan, and H. N. Nagaraja. *A first course in order statistics*, volume 54. Siam, 1992.
- [2] J. Avérous, C. Genest, and S. C. Kochar. On the dependence structure of order statistics. *Journal of Multivariate Analysis*, 94(1):159–171, 2005.
- [3] P. J. Bickel. Some contributions to the theory of order statistics. In *Proc. Fifth Berkeley Sympos. Math. Statist. and Probability (Berkeley, Calif., 1965/66), Vol. I: Statistics*, pages 575–591. Univ. California Press, Berkeley, Calif., 1967.
- [4] P. J. Boland, M. Hollander, K. Joag-Dev, and S. Kochar. Bivariate dependence properties of order statistics. *Journal of Multivariate Analysis*, 56(1):75–89, 1996.
- [5] J. Borwein, A. Lewis, and R. Nussbaum. Entropy minimization, *DAD* problems, and doubly stochastic kernels. *Journal of Functional Analysis*, 123(2):264 – 307, 1994. ISSN 0022-1236. doi: <http://dx.doi.org/10.1006/jfan.1994.1089>. URL <http://www.sciencedirect.com/science/article/pii/S0022123684710895>.
- [6] C. Butucea, J.-F. Delmas, A. Dutfoy, and R. Fischer. Maximum entropy copula with given diagonal section. *Journal of Multivariate Analysis*, 137:61 – 81, 2015. ISSN 0047-259X. doi: <http://dx.doi.org/10.1016/j.jmva.2015.01.003>. URL <http://www.sciencedirect.com/science/article/pii/S0047259X15000081>.
- [7] H. A. David and H. N. Nagaraja. *Order statistics*. Wiley Online Library, 1970.
- [8] B. V. de Melo Mendes and M. A. Sanfins. The limiting copula of the two largest order statistics of independent and identically distributed samples. *Brazilian Journal of Probability and Statistics*, 21:85–101, 2007.
- [9] D. Dubhashi and O. Häggström. A note on conditioning and stochastic domination for order statistics. *J. Appl. Probab.*, 45(2):575–579, 2008. ISSN 0021-9002. doi: 10.1239/jap/1214950369. URL <http://dx.doi.org/10.1239/jap/1214950369>.
- [10] T. Hu and H. Chen. Dependence properties of order statistics. *Journal of Statistical Planning and Inference*, 138(7):2214–2222, 2008.
- [11] P. Jaworski. On copulas and their diagonals. *Information Sciences*, 179(17):2863 – 2871, 2009. ISSN 0020-0255. doi: 10.1016/j.ins.2008.09.006. URL <http://www.sciencedirect.com/science/article/pii/S0020025508003836>.
- [12] P. Jaworski and T. Rychlik. On distributions of order statistics for absolutely continuous copulas with applications to reliability. *Kybernetika*, 44(6):757–776, 2008.
- [13] S. Kim and H. David. On the dependence structure of order statistics and concomitants of order statistics. *Journal of statistical planning and inference*, 24(3):363–368, 1990.
- [14] R. Lebrun and A. Dutfoy. Copulas for order statistics with prescribed margins. *Journal of Multivariate Analysis*, 128:120–133, 2014.
- [15] J. Navarro and N. Balakrishnan. Study of some measures of dependence between order statistics and systems. *Journal of Multivariate Analysis*, 101(1):52–67, 2010.
- [16] J. Navarro and F. Spizzichino. On the relationships between copulas of order statistics and marginal distributions. *Statist. Probab. Lett.*, 80(5-6):473–479, 2010. ISSN 0167-7152. doi: 10.1016/j.spl.2009.11.025. URL <http://dx.doi.org/10.1016/j.spl.2009.11.025>.
- [17] L. Rüschendorf and W. Thomsen. Note on the Schrödinger equation and *I*-projections. *Statist. Probab. Lett.*, 17(5):369–375, 1993. ISSN 0167-7152. doi: 10.1016/0167-7152(93)90257-J. URL [http://dx.doi.org/10.1016/0167-7152\(93\)90257-J](http://dx.doi.org/10.1016/0167-7152(93)90257-J).
- [18] V. Schmitz. Revealing the dependence structure between $X_{(1)}$ and $X_{(n)}$. *Journal of statistical planning and inference*, 123(1):41–47, 2004.

- [19] N. Zhao and W. T. Lin. A copula entropy approach to correlation measurement at the country level. *Applied Mathematics and Computation*, 218(2):628 – 642, 2011. ISSN 0096-3003. doi: 10.1016/j.amc.2011.05.115. URL <http://www.sciencedirect.com/science/article/pii/S0096300311007983>.

CRISTINA BUTUCEA, UNIVERSITÉ PARIS-EST, LAMA (UPE-MLV), 77455 MARNE LA VALLÉE, FRANCE.
E-mail address: `cristina.butucea@univ-mlv.fr`

JEAN-FRANÇOIS DELMAS, UNIVERSITÉ PARIS-EST, CERMICS (ENPC), 77455 MARNE LA VALLÉE, FRANCE.
E-mail address: `delmas@cermics.enpc.fr`

ANNE DUTFOY, EDF RESEARCH & DEVELOPMENT, INDUSTRIAL RISK MANAGEMENT DEPARTMENT, 92141 CLAMART CEDEX, FRANCE.
E-mail address: `anne.dutfoy@edf.fr`

RICHARD FISCHER, UNIVERSITÉ PARIS-EST, CERMICS (ENPC), 77455 MARNE LA VALLÉE, FRANCE, EDF RESEARCH & DEVELOPMENT, INDUSTRIAL RISK MANAGEMENT DEPARTMENT, 92141 CLAMART CEDEX, FRANCE.
E-mail address: `fischerr@cermics.enpc.fr`